

## DAFTAR PUSTAKA

- [1] C. P. 'na B. and N. M. ~ 'ia C. Guzman V ' azquez, "Stoichiometric hydroxyapatite obtained by precipitation and sol gel processes," 2004.
- [2] M. Fitriawan, "Sintesis Hidroksiapatit Berbahan Dasar Tulang Sapi dengan Metode Wet Process sebagai Material Pengganti Graft Sintesis Hidroksiapatit Berbahan Dasar Tulang Sapi dengan Metode Preti ...," no. April, 2016, doi: 10.13140/RG.2.1.4571.3449.
- [3] M. Tomi, "Pengaruh Jenis dan Jumlah Bola Penggiling terhadap Karakteristik Fisik Serbuk Cangkang Telur Ayam dengan Penggilingan Ball Mill," 2016, [Online]. Available: <http://scholar.unand.ac.id/18031/>
- [4] K. U. Henggu, B. Ibrahim, and P. Suptijah, "Hidroksiapatit dari cangkang sotong sebagai sediaan biomaterial perancah tulang," *Jphpi*, vol. 22, no. 1, pp. 1–13, 2019.
- [5] R. Purwo Ningsih, N. Wahyuni, and L. Destiarti, "Sintesis Hidroksiapatit Dari Cangkang Kerang Kepah (Polymesoda erosa) Dengan Variasi Waktu Pengadukan."
- [6] G. Dwi Putra, "Pembuatan dan Karakterisasi Serbuk Nanob Biokeramik Dari Cangkang Telur Ayam Ras yang Dilakukan Dengan Proses Ball Mill dan Pemanasan Sela," Universitas Andalas, 2016.
- [7] M. A. N. R. Bambang, "Material Biokeramik Berbasis Hidroksiapatit Tulang Ikan Tuna," *Jurnal Departemen Teknologi Hasil Pertanian Fakultas Perikanan dan Ilmu Kelautan Institut Pertanian Bogor*, pp. 119–132, 2013.
- [8] R. J. Reis, "On some new loricarides," pp. 277–278 .
- [9] R. Covain, by Claude Weber, raphaël Covain, and S. FiSch-Muller, "Identity of *Hypostomus plecostomus* (Linnaeus, 1758), with an overview of *Hypostomus* species from the Guianas (Teleostei: Siluriformes: Loricariidae)," 2012. [Online]. Available: <https://www.researchgate.net/publication/237021825>
- [10] Silva et al, "Nutritional composition and sensory evaluation of *Plecostomus plecostomus* (Hypostomidae) from the Amazon region," 2017.

- [11] M. Boutinguiza, J. Pou, R. Comesaña, F. Lusquiños, A. De Carlos, and B. León, “Biological hydroxyapatite obtained from fish bones,” *Materials Science and Engineering C*, vol. 32, no. 3, pp. 478–486, Apr. 2012, doi: 10.1016/j.msec.2011.11.021.
- [12] R. Othman and A. Zakaria, “OPTIMISATION OF MILLING PARAMETERS DURING MECHANICAL ACTIVATION FOR DIRECT SYNTHESIS OF HYDROXYAPATITE.”
- [13] Y. Kaya, V. Kobya, A. Mardani, N. Mardani, and H. E. Beytekin, “Effect of Grinding Conditions on Clinker Grinding Efficiency: Ball Size, Mill Rotation Speed, and Feed Rate,” *Buildings*, vol. 14, no. 8, Aug. 2024, doi: 10.3390/buildings14082356.
- [14] A. Fadli, S. R. Yenti, A. Prabowo, S. P. Aledya, R. Fadila, and R. Susanto, “The Synthesis of Hydroxyapatite Powder Using Shaker Mill Technique,” *Journal of Southwest Jiaotong University*, vol. 56, no. 2, pp. 22–31, Apr. 2021, doi: 10.35741/issn.0258-2724.56.2.3.
- [15] A. dkk Hammood, “Effect of calcination temperature on characterization of natural hydroxyapatite prepared from carp fish bones,” vol. 1, no. 436, 2019.
- [16] Piccirillo C dkk., “Hydroxyapatite and chloroapatite derived from sardine by- products ,” 2017.
- [17] Pal Anindya dkk, “Synthesis of hydroxyapatite from Lates Calcarifer fish bone for Biomedical Applications,” vol. 203, no. 89–92, 2017.
- [18] Marlina Ana dkk, “Sythesis and Characterization of Hydroxiapatite from fish Bone Waste ,” 2015.
- [19] S. M. S. Hariani P, “Effect of Sintering on the Mechanical Properties of Hydroxyapatite From Fish Bone (Pangasius Hypophthalmus,” 2019.
- [20] E. Fibryanto and H. Amanda, “Characterization of nano-hydroxyapatite–collagen and epigallocatechin-3-gallate (EGCG) composites by scanning electron microscopy-energy dispersive spectroscopy (SEM-EDS), X-ray diffraction (XRD), and Fourier transform infrared (FTIR) spectroscopy,” *Scientific Dental Journal*, vol. 6, no. 2, p. 80, 2022, doi: 10.4103/sdj.sdj\_4\_22.