

DAFTAR PUSTAKA

1. Ye Y, Zhao Y, Han T, Zhang X, Miao H, Qin B, et al. Accuracy of axial length, keratometry, and refractive measurement with Myopia Master in children with ametropia. *BMC Ophthalmol.* 2022;22(1):1–7.
2. Holden BA, Fricke TR, Wilson DA, Jong M, Naidoo KS, Sankaridurg P, et al. Global prevalence of myopia and high myopia and temporal trends from 2000 through 2050. *Ophthalmology.* 2016;123(5):1036–42.
3. Brodie SE, Ang M, Irsch K, Jackson ML, Mauger TF, Oostra T et al. Optics of the human eye. In: *Clinical Optics and Vision Rehabilitation.* San Fransisco: American Academy of Ophthalmology; 2024. p. 143–63.
4. Jiang F, Wang D, Yin Q, He M, Li Z. Longitudinal Changes in Axial Length and Spherical Equivalent in Children and Adolescents With High Myopia. *Investig Ophthalmol Vis Sci.* 2023;64(12):1–4.
5. Chung HJ, Park CK. The Distinct Biometric Features of High Myopia Compared to Moderate Myopia. *Curr Eye Res.* 2016;41(12):1580–3.
6. Ohno-matsui K, Wu P chang, Yamashiro K, Vutipongsatorn K, Fang Y, Ming C, et al. IMI Pathologic Myopia. *Investig Ophthalmol Vis Sci.* 2021;62:1–36.
7. Morgan IG, Ohno-Matsui K, Saw SM. Myopia. In: *The Lancet.* 2012. p. 1739–48.
8. Jiang Y, Wang D, Han X, Liao C, Li Z, Scheetz J, et al. Visual impairment in highly myopic eyes: The ZOC-BHVI high myopia cohort study. *Clin Exp Ophthalmol.* 2020;48(6):783–92.
9. Zhao X, Ding X, Lyu C, Li S, Liu B, Li T, et al. Morphological characteristics and visual acuity of highly myopic eyes with different severities of myopic maculopathy. *Retina.* 2020;40(3):461–7.
10. Liu HH, Xu L, Wang YX, Wang S, You QS, Jonas JB. Prevalence and progression of myopic retinopathy in Chinese adults: The Beijing eye study. *Ophthalmology.* 2010;117(9):1763–8.
11. Tideman JW, Snabel MCC, Tedja MS, Van Rijn GA, Wong KT, Kuijpers RAM, et al. Association of axial length with risk of uncorrectable visual impairment for europeans with myopia. *JAMA Ophthalmol.* 2016;134(12):1355–63.
12. Wang Y, Wang D, Yin Q, Li J, Li Z, He M. Choroidal thickness and visual acuity in high myopia without myopic maculopathy: Insights from a chinese population study. *Transl Vis Sci Technol.* 2024;13(11):1–10.
13. Troilo D, Smith EL, Nickla DL, Ashby R, Tkatchenko A V., Ostrin LA, et al. IMI – Report on experimental models of emmetropization and myopia. *Investig Ophthalmol Vis Sci.* 2019;60(3):M31–88.
14. Gupta P, Thakku SG, Saw SM, Tan M, Lim E, Tan M, et al. Characterization of choroidal morphologic and vascular features in young men with high myopia using spectral-domain optical coherence tomography. *Am J Ophthalmol.* 2017;177:27–33.
15. Fujiwara T, Imamura Y, Margolis R, Slakter JS, Spaide RF. Enhanced depth imaging optical coherence tomography of the choroid in highly myopic eyes. *Am J Ophthalmol.* 2009;148.
16. Ye J, Shen M, Huang S, Fan Y, Yao A, Pan C, et al. Visual acuity in pathological myopia is correlated with the photoreceptor myoid and ellipsoid zone thickness and affected by choroid thickness. *Investig Ophthalmol Vis Sci.* 2019;60(5):1714–23.
17. Read SA, Collins MJ, Vincent SJ, Alonso-Caneiro D. Choroidal thickness in myopic and nonmyopic children assessed with enhanced depth imaging optical coherence tomography. *Investig Ophthalmol Vis Sci.* 2013;54(12):7578–86.

18. Zhou LX, Shao L, Xu L, Wei W Bin, Wang YX, You QS. The relationship between scleral staphyloma and choroidal thinning in highly myopic eyes: The Beijing Eye Study. *Sci Rep.* 2017;7(1):3–8.
19. Chakraborty R, Read SA VS. Understanding Myopia: Patogenesis and Mechanisms. In: *Updates on Myopia: A Clinical Perspective.* Singapore: Springer; 2020. p. 65–94.
20. Howlett MHC, McFadden SA. Form-deprivation myopia in the guinea pig (*Cavia porcellus*). *Vision Res.* 2006;46(1–2):267–83.
21. Chalam K V., Sambhav K. Choroidal thickness measured with swept source optical coherence tomography in posterior staphyloma strongly correlates with axial length and visual acuity. *Int J Retin Vit.* 2019;5(1):1–9.
22. Sun B, Ye Y, Zhao J, Zhou X, Niu L. Comparison of axial length and anterior segment parameters of patients with myopia measured using 2 fourier-domain optical coherent biometry devices. *BMC Ophthalmol.* 2024;24(1):1–8.
23. Zhang Z, Mu J, Wei J, Geng H, Liu C, Yi W, et al. Correlation between refractive errors and ocular biometric parameters in children and adolescents: a systematic review and meta-analysis. *BMC Ophthalmol.* 2023;23(1):1–24.
24. Sabur H, Takes O. Agreement of axial length and anterior segment parameters measured with the MYAH device compared to Pentacam AXL and IOLMaster 700 in myopic children. *Int Ophthalmol.* 2023;43(2):475–82.
25. Spaide RF, Koizumi H, Pozzoni MC. Enhanced depth imaging spectral-domain optical coherence tomography. *Am J Ophthalmol.* 2008;146.
26. Benjamin WJ. Refractive Status of The Eye. In: *Borish's Clinical Refraction.* Second Edi. Philadelphia: Elsevier Inc; 2007.
27. Dolgin E. The myopia boom. *Nature.* 2015;519(7543):276–8.
28. Jung SK, Lee JH, Kakizaki H, Jee D. Prevalence of myopia and its association with body stature and educational level in 19-year-old male conscripts in Seoul, South Korea. *Investig Ophthalmol Vis Sci.* 2012;53(9):5579–83.
29. Wong TY, Ferreira A, Hughes R, Carter G, Mitchell P. Epidemiology and disease burden of pathologic myopia and myopic choroidal neovascularization: an evidence-based systematic review. *Am J Ophthalmol.* 2014;157.
30. Chen M, Wu A, Zhang L, Wang W, Chen X, Yu X, et al. The increasing prevalence of myopia and high myopia among high school students in Fenghua city, eastern China: A 15-year population-based survey. *BMC Ophthalmol.* 2018;18(1):1–10.
31. Wu JF, Bi HS, Wang SM, Hu YY, Wu H, Sun W, et al. Refractive error, visual acuity and causes of vision loss in children in Shandong, China. The Shandong children eye study. *PLoS One.* 2013;8(12).
32. Koh V, Yang A, Saw SM, Chan YH, Lin ST, Tan MMH, et al. Differences in prevalence of refractive errors in young asian males in singapore between 1996-1997 and 2009-2010. *Ophthalmic Epidemiol.* 2014;21(4):247–55.
33. Ruterjerg J, Ilmer M, Recio A, Coleman M, Vykoukal J, Alt E, et al. Myopia in young adults is inversely related to an objective marker of ocular sun exposure: The Western Australian raine cohort study. *Front Neuroendocrinol.* 2014;35(3):320–30.
34. French AN, Morgan IG, Burlutsky G, Mitchell P, Rose KA. Prevalence and 5- to 6-year incidence and progression of myopia and hyperopia in Australian schoolchildren. *Ophthalmology.* 2013;120(7):1482–91.
35. Jacobsen N, Jensen H, Goldschmidt E. Prevalence of myopia in Danish conscripts. *Acta Ophthalmol Scand.* 2007;85(2):165–70.
36. Midelfart A, Kinge B, Midelfart S, Lydersen S. Prevalence of refractive errors in young and middle-aged adults in Norway. *Acta Ophthalmol Scand.* 2002;80(5):501–5.

37. Sun J, Zhou J, Zhao P, Lian J, Zhu H, Zhou Y, et al. High prevalence of myopia and high myopia in 5060 Chinese University students in Shanghai. *Investig Ophthalmol Vis Sci.* 2012;53(12):7504–9.
38. Rose KA, Morgan IG, Smith W, Burlutsky G, Mitchell P, Saw SM. Myopia, lifestyle, and schooling in students of Chinese ethnicity in Singapore and Sydney. *Arch Ophthalmol.* 2008;126(4):527–30.
39. Saw SM, Gazzard G, Koh D, Farook M, Widjaja D, Lee J, et al. Prevalence rates of refractive errors in Sumatra, Indonesia. *Investig Ophthalmol Vis Sci.* 2002;43(10):3174–80.
40. Mahayana IT, Indrawati SG, Pawiroranu S. The prevalence of uncorrected refractive error in urban, suburban, exurban and rural primary school children in Indonesian population. *Int J Ophthalmol.* 2017;10(11):1771–6.
41. Moderiano D, Do M, Hobbs S, Lam V, Sarin S, Alonso-Caneiro D, et al. Influence of the time of day on axial length and choroidal thickness changes to hyperopic and myopic defocus in human eyes. *Exp Eye Res.* 2019;182(April 2018):125–36.
42. Nishijima David L; Wisner, David H; Holmes, James F DKS, Maxson & Mitchell. The dynamic sclera: Extracellular matrix remodeling in normal ocular growth and myopia development. *Physiol Behav.* 2016;176(25):139–48.
43. dan Bree D levy. Dopamine signaling and myopia development: What are the key challenges. *Physiol Behav.* 2019;176(3):139–48.
44. Xiang ZY, Zou HD. Recent Epidemiology Study Data of Myopia. *J Ophthalmol.* 2020;2020.
45. Mirshahi A, Ponto KA, Hoehn R, Zwiener I, Zeller T, Lackner K, et al. Myopia and level of education. *Am Acad Ophthalmol Ophthalmol J Ophthalmol.* 2014;11(1):2047–52.
46. Jones LA, Sinnott LT, Mutti DO, Mitchell GL, Melvin L, Zadnik K. Parental aistory of myopia, sports and outdoor activities, and future myopia. *Nature.* 2010;23(1):1–7.
47. Rose KA, Morgan IG, Ip J, Kifley A, Huynh S, Smith W, et al. Outdoor activity reduces the prevalence of myopia in children. *Ophthalmology.* 2008;115(8):1279–85.
48. Ohno-Matsui K, Lai TYY, Lai CC, Cheung CMG. Updates of pathologic myopia. *Prog Retin Eye Res.* 2016;52:156–87.
49. Ohno-Matsui K, Kawasaki R, Jonas JB, Cheung CMG, Saw SM, Verhoeven VJM, et al. International photographic classification and grading system for myopic maculopathy. *Am J Ophthalmol.* 2015;159(5):877-883.e7.
50. Verkicharla PK, Ohno-Matsui K, Saw SM. Current and predicted demographics of high myopia and an update of its associated pathological changes. *Ophthalmic Physiol Opt.* 2015;35(5):465–75.
51. Bhardwaj V, Rajeshbhai GP. Axial length, anterior chamber depth-a study in different age groups and refractive errors. *J Clin Diagnostic Res.* 2013;7(10):2211–2.
52. Dogan M, Elgin U, Sen E, Tekin K, Yilmazbas P. Comparison of anterior segment parameters and axial lengths of myopic, emmetropic, and hyperopic children. *Int Ophthalmol.* 2019;39(2):335–40.
53. Brar VS, Couser NL, Dhalla MS, Goldman DJ, Kang KB, Richards NQ SJ. *The Eye. In: Fundamentals and Principles of Ophthalmology.* San Fransisco: American Academy of Ophthalmology; p. 79–82.
54. Kim SJ, Fawzi A, Kovach JL, Petel S, Recchia FM, Sobrin L, Sun JK WC. Basic anatomy. *In: Retina and Vitreous.* San Fransisco: American Academy of Ophthalmology; p. 17–9.
55. Wei W Bin, Xu L, Jonas JB, Shao L, Du KF, Wang S, et al. Subfoveal choroidal

- thickness : The Beijing. 2012;175–80.
56. Read SA, Alonso-Caneiro D, Vincent SJ, Collins MJ. Longitudinal changes in choroidal thickness and eye growth in childhood. *Investig Ophthalmol Vis Sci.* 2015;56(5):3103–10.
 57. Wei W Bin, Xu L, Jonas JB, Shao L, Du KF, Wang S, et al. Subfoveal choroidal thickness: The Beijing eye study. *Ophthalmology.* 2013;120(1):175–80.
 58. Read SA, Collins MJ, Vincent SJ. Light exposure and eye growth in childhood. *Investig Ophthalmol Vis Sci.* 2015;56(11):6779–87.
 59. Kong M, Choi DY, Han G, Song YM, Park SY, Sung J, et al. Measurable range of subfoveal choroidal thickness with conventional spectral domain optical coherence tomography. *Transl Vis Sci Technol.* 2018;7(5).
 60. Entezari M, Karimi S, Ramezani A, Nikkhah H, Fekri Y, Kheiri B. Choroidal thickness in healthy subjects. *J Ophthalmic Vis Res.* 2018;13(1):39–43.
 61. Yavuzer K, Bozkurt B, Ozturk BT, Okudan S. The effect of age on subfoveal choroidal thickness in healthy subjects. *East J Med.* 2021;26(4):519–25.
 62. Liu B, Wang Y, Li T, Lin Y, Ma W, Chen X, et al. Correlation of subfoveal choroidal thickness with axial length, refractive error, and age in adult highly myopic eyes. *BMC Ophthalmol.* 2018;18(1):1–5.
 63. Gupta P, Cheung CY, Saw SM, Koh V, Tan M, Yang A, et al. Choroidal thickness does not predict visual acuity in young high myopes. *Acta Ophthalmol.* 2016;94.
 64. Gupta P, Saw SM, Cheung CY, Girard MJ, Mari JM, Bhargava M, et al. Choroidal thickness and high myopia: a case-control study of young Chinese men in Singapore. *Acta Ophthalmol.* 2015;93.
 65. Al-abri M, Al-aliyani A, Mal W, Holla A, Al-rawahi A, Al-marhoon M. Choroidal thickness profile in healthy eyes of adult omani population : A cross-sectional , descriptive clinical study. 2022;13(1000935).
 66. Sun Y, Xu F, Zhang T, Liu M, Wang D, Chen Y, et al. Orthokeratology to control myopia progression : A meta-analysis. 2015;1–9.
 67. Wu PC, Chen CT, Lin KK, Sun CC, Kuo CN, Huang HM, et al. Myopia prevention and outdoor light intensity in a school-based cluster randomized trial. *Ophthalmology.* 2018;125(8):1239–50.
 68. French AN, Ashby RS, Morgan IG, Rose KA. Time outdoors and the prevention of myopia. *Exp Eye Res.* 2013;(May):1–11.
 69. Mccarthy CS, Megaw P, Devadas M, Morgan IG. Dopaminergic agents affect the ability of brief periods of normal vision to prevent form-deprivation myopia. 2007;84:100–7.
 70. Wu LJ, You QS, Duan JL, Luo YX, Liu LJ, Li X, et al. Prevalence and associated factors of myopia in high-school students in Beijing. *PLoS One.* 2015;10.
 71. Hashemi H, Rezvan F, Beiranvand A, Papi O ali. Prevalence of refractive errors among high school students in Western Iran. 2014;9(2):232–9.
 72. Margolis RON, Spaide RF. A pilot study of enhanced depth imaging optical coherence tomography of the choroid in normal eyes. 2009;811–5.
 73. Flores-moreno I, Lugo F, Duker JAYS. The relationship between axial length and choroidal thickness in eyes with high myopia . 2013;
 74. Yin G, Wang YX, Zheng ZY, Yang H, Xu L, Jonas JB, et al. Ocular axial length and its associations in Chinese : The Beijing eye study. 2012;7(8).
 75. Purpose K, Journal K. Relations between age, weight, refractive error and eye shape by computerized tomography in children. 2006;(January 2005):7–10.
 76. Cunha CM, Marchezine G, Cunha JT, Morais G, Lima B De, Martins I, et al. Refractive

- Error and Ocular Biometry Among Young adults From Cuiabá , 2023;1–5.
77. Yamashita T, Yamashita T, Shirasawa M, Arimura N. Repeatability and reproducibility of subfoveal choroidal thickness in normal eyes of japanese using different SD-OCT devices. 2012;53(3):1102–7.
 78. Takahashi A, Ito Y, Iguchi Y, Yasuma TR, Ishikawa K, Terasaki H. Axial length increases and related changes in highly myopic normal eyes with myopic complications in fellow eyes.
 79. Gupta P, Thakku SG, Saw S mei, Tan M, Lim E, Tan M, et al. Detailed phenotyping and characterization of choroidal morphological and vascular features in young men with high myopia using Spectral-Domain Optical Coherence Tomography. 2017;
 80. Touhami S, Philippakis E, Mrejen S, Couturier A, Casteran C, Levent P, et al. Topographic variations of choroidal thickness in healthy eyes on swept-source optical coherence. *Investig Ophthalmol Vis Sci.* 2019;61(3):1–11.
 81. Muhiddin HS, Mayasari AR, Umar BT, Sirajuddin J, Patellongi I, Islam IC, et al. Choroidal thickness in correlation with axial length and myopia degree. 2022;1–9.
 82. Zhao M, Kwok A, Lam C, Tin M, Ying C. Hemodynamic and morphological changes of the central retinal artery in myopic eyes. *Sci Rep.* 2022;(0123456789):1–9.
 83. Kong H. Choroidal thickness and visual acuity in highly myopic eyes.
 84. Chui TYP, Yap MKH, Chan HHL, Thibos LN. Retinal stretching limits peripheral visual acuity in myopia. 2005;45:593–605.
 85. Park UC, Yoon CK, Bae K, Lee EK. Association of retinal sensitivity with optical coherence tomography microstructure in highly myopic patients. 2022;(February 2020).
 86. Zhou Z, Xiong P, Sun J, Wang Y, Wang J. Effects of posterior staphyloma on choroidal structure in myopic adults : a retrospective study. *BMC Ophthalmol.* 2023;1–10.
 87. Li Y, Li H, Rui X, Wang Y, Zhu S, Huang M, et al. Choroidal vascular changes in early - stage myopic maculopathy from deep learning choroidal analysis : a hospital - based SS - OCT study. 2024;1–13.
 88. Article O, Teberik K, Kaya M. Retinal and choroidal thickness in patients with high myopia without maculopathy. 2017;33(6):1438–43.
 89. Summers JA, Shelton S, Norton TT. The sclera and myopia. 2006;82:185–200.
 90. Agrawal R, Gupta P, Tan K anne, Ming C, Cheung G. Choroidal vascularity index as a measure of vascular status of the choroid : Measurements in healthy eyes from a population-based study. *Nat Publ Gr.* 2016;(February):1–9.
 91. Ruiz-Medrano J, Flores-Moreno I, Pena-Garcia P, Montero JA, Duker JS, Ruiz-Moreno JM. Macular choroidal thickness profile in a healthy population measured by swept-source optical coherence tomography. *Invest Ophthalmol Vis Sci.* 2014;55.
 92. Gunkel M, Scho J, Kaupp UB, Al-amoudi A, Kaupp UB, Alkhalidi W, et al. Higher-order architecture of rhodopsin in intact photoreceptors and its implication for phototransduction kinetics article higher-order architecture of rhodopsin in intact photoreceptors and its implication for phototransduction kinetics. 2015;628–38.
 93. Wong CW, Phua V, Lee SY, Wong TY, Cheung CM. Is choroidal or scleral thickness related to myopic macular degeneration? *Invest Ophthalmol Vis Sci.* 2017;58.
 94. Feldkaemper M, Schaeffel F. An updated view on the role of dopamine in myopia. *Exp Eye Res.* 2013;114:106–19.
 95. Chia A, Lu Q shu, Tan D. Five-year clinical trial on atropine for the treatment of myopia 2: Myopia control with atropine 0 . 01 % eyedrops. 2015;1–9.
 96. Fang Y, Du R, Nagaoka N, Yokoi T, Shinohara K, Xu X, et al. OCT-based diagnostic criteria for different stages of myopic maculopathy. 2019;1–15.

97. Flitcroft DI. The complex interactions of retinal , optical and environmental factors in myopia aetiology. *Prog Retin Eye Res.* 2012;31(6):622–60.
98. Jonas JB, Ang M, Cho P, Guggenheim JA, He MG, Jong M, et al. IMI prevention of myopia and its progression. 2021;(Imi):8–10.
99. Song Y yan, Wang H, Wang B song, Qi H, Rong Z xing, Chen H zhuan. Atropine in ameliorating the progression of myopia in children with mild to moderate myopia : A meta-analysis of controlled clinical trials. 2011;27(4).
100. Randomized A, Trial P controlled, Yam JC, Edin F, Jiang Y, Tang SM, et al. Low-concentration atropine for myopia progression (LAMP) study myopia control. 2018;1–12.
101. Sankaridurg P, Holden B, Smith E, Naduvilath T, Chen X, de la Jara PL, et al. Decrease in rate of myopia progression with a contact lens designed to reduce relative peripheral hyperopia: One-year results. *Investig Ophthalmol Vis Sci.* 2011;52(13):9362–7.

