

DAFTAR PUSTAKA

- Agarwal, V. M., Rastogi, N., & Raju, S. V. S. (2007). Impact of predatory ants on two lepidopteran insect pests in Indian cauliflower agroecosystems. *Journal of Applied Entomology*, 131(7), 493–500. <https://doi.org/10.1111/j.1439-0418.2007.01199.x>
- Alves, L. F. ., Bellon, P. P., Rheinheimer, A. R., & Pietrowski, V. (2012). First record of *Beauveria bassiana* (Hyphomycetes: Moniliales) on adults of cassava lace bug *Vatiga manihotae* (Drake) (Hemiptera: Tingidae) in Brazil. *Arquivos do Instituto Biológico*, 79(2), 309–311. <https://doi.org/10.1590/s1808-16572012000200024>
- Ayudya, D. R., Herlinda, S., & Suwandi, S. (2019). Insecticidal activity of culture filtrates from liquid medium of *Beauveria bassiana* isolates from South Sumatra (Indonesia) wetland soil against larvae of *Spodoptera litura*. *Biodiversitas*, 20(8), 2101–2109. <https://doi.org/10.13057/biodiv/d200802>
- BPS Kabupaten Lima Puluh Kota. (2024). Lima Puluh Kota Dalam Angka 2024 (B. P. S. K. L. P. Kota/ (Ed.)). Badan Pusat Statistik Kabupaten Lima Puluh Kota/
- BPS Sumatera Barat. (2023). Sumatera Barat Dalam Angka 2024. In Berita Resmi Badan Pusat Statistik (Vol. 54).
- Basset, Y., Missa, O., Alonso, A., Miller, S. E., Curletti, G., De Meyer, M., Eardley, C., Lewis, O. T., Mansell, M. W., Novotny, V., & Wagner, T. (2008). Monitoring arthropods in a tropical landscape: Relative effects of sampling methods and habitat types on trap catches. *Journal of Insect Conservation*, 12(2), 143–159. <https://doi.org/10.1007/s10841-007-9130-5>
- Begg, G. S., Cook, S. M., Dye, R., Ferrante, M., Franey, P., Lavigne, C., Lövei, G. L., Mansion-Vaquie, A., Pell, J. K., Petit, S., Quesada, N., Ricci, B., Wratten, S. D., & Birch, A. N. E. (2017). A functional overview of conservation biological control. *Crop Protection*, 97, 145–158. <https://doi.org/10.1016/j.cropro.2016.11.008>
- Bellon, P. P., Oliveira, H. N. De, Loureiro, E. D. S., Renato, D., Santana, S., Otsubo, A. A., & Mota, T. A. (2017). Populational fluctuation of lace bug in cassava. 1–6. <https://doi.org/10.1590/1808-1657000602015>
- Bellon PP, da Silva Wengrat APG, Kassab SO, Pietrowski V, L. E. (2012). Occurrence of Lace bug *Vatiga illudens* and *Vatiga manihotae* (Hemiptera : Tingidae) in Mato Grosso do Sul , midwestern Brazil. *Annals of the Brazilian Academy of Sciences*, 84, 703–705.
- Bellotti, A. C., Arias V., B., Vargas H., O., & Peña, J. E. (2002). Pérdidas en rendimiento del cultivo de Yuca causadas por insectos y ácaros. La yuca en Tercer Milenio: Sistemas modernos de producción, procesamiento, utilización y comercialización. Concorcio Latinoamericano para la Investigación y el Desarrollo de la yuca; Proyecto IP-3 de Mejoramiento de Yuca, Cali, CO. Publicación CIAT no. 327, 47, 204–219.

- Bellotti, A. C., Melo, E. L., Arias, B., Herrera, C. J., Hernández, M. del P., Holguín, C. M., Guerrero, J. M., & Trujillo, H. (2005). Biological control in the neotropics: A selective review with emphasis on cassava. Second International Symposium on Biological Control of Arthropods; September 12-16, 2005, Davos, Switzerland, 1(December), 206–227.
- Bellotti, A. C., Smith, L., & Lapointe, S. L. (1999). Recent advances in cassava pest management. *Annual Review of Entomology*, 44, 343–370. <https://doi.org/10.1146/annurev.ento.44.1.343>
- Bellotti, A., Campo, B. V. H., & Hyman, G. (2012). Cassava Production and Pest Management: Present and Potential Threats in a Changing Environment. In *Tropical Plant Biology* (Vol. 5, Nomor 1, hal. 39–72). <https://doi.org/10.1007/s12042-011-9091-4>
- Bianchi, F. J. J. A., Booij, C. J. H., & Tschamntke, T. (2006). Sustainable pest regulation in agricultural landscapes: A review on landscape composition, biodiversity and natural pest control. *Proceedings of the Royal Society B: Biological Sciences*, 273(1595), 1715–1727. <https://doi.org/10.1098/rspb.2006.3530>
- Chalil, D. (2003). *Agribisnis Ubi Kayu di Propinsi Sumatera Utara*. Jurusan Sosial Ekonomi Pertanian, Fakultas Pertanian, Universitas Sumatera Utara. Medan.
- Choate, B., & Drummond, F. (2011). Ants as biological control agents in agricultural cropping systems. *Terrestrial Arthropod Reviews*, 4(2), 157–180. <https://doi.org/10.1163/187498311X571979>
- Cho, G., Kim, W., Kwon, Y. S., & Lee, S. (2020). Check list of lace bugs (Hemiptera: Tingidae) of Korea: Tingidae check list of Korea. *Journal of Asia-Pacific Entomology*, 23(3), 736–745. <https://doi.org/10.1016/j.aspen.2020.04.005>
- Duelli, P., Obrist, M. K., & Schmatz, D. R. (1999). Biodiversity evaluation in agricultural landscapes: Above-ground insects. *Agriculture, Ecosystems and Environment*, 74(1–3), 33–64. [https://doi.org/10.1016/S0167-8809\(99\)00029-8](https://doi.org/10.1016/S0167-8809(99)00029-8)
- Du, S., & Yao, Y. (2018). A new genus and species of Tingidae (Heteroptera : Cimicomorpha) from Myanmar , with the analysis of the evolution of hood , carinae and paranota. *Zoological Systematics*, 43(June), 283–293. <https://doi.org/10.11865/zs.201828>
- Erdi S., R. (2016). Kelimpahan musuh alami (predator) pada tanaman jagung di desa saree kecamatan lembah seulawah kabupaten aceh besar. IV(2), 10–18.
- Fialho, J. F. E. A., Silva, S. V. de P.-M. M. S., & Junqueira, N. T. V. (2009). Danos Causados Por Percevejo-De-Renda Na Produção De Parte Aérea E Raízes De Mandioca Economic Damage Caused By Lacebug Upon Cassava Root And. *Scientia Agraria, Curitiba*, 1125, 151–155.
- Froeschner, R. C. (1993). The neotropical lace bugs of the genus *Vatiga* (Heteroptera : Tingidae), pests of cassava : new synonymies and key to species. *Proceedings of the Entomological Society of Washington*, 95(3), 457–462.

- Guidoti, M., Montemayor, S. I., & Guilbert, É. (2015). True Bugs (Heteroptera) Of The Neotropics. *True Bugs (Heteroptera) Of The Neotropics*, April 2016, 1–862. <https://doi.org/10.1007/978-94-017-9861-7>
- Gurr, G. M., Lu, Z., Zheng, X., Xu, H., Zhu, P., Chen, G., Yao, X., Cheng, J., Zhu, Z., Catindig, J. L., Villareal, S., Van Chien, H., Cuong, L. Q., Channoo, C., Chengwattana, N., Lan, L. P., Hai, L. H., Chaiwong, J., Nicol, H. I., ... Heong, K. L. (2016). Multi-country evidence that crop diversification promotes ecological intensification of agriculture. *Nature Plants*, 2(3), 16014. <https://doi.org/10.1038/nplants.2016.14>
- Gurr, G. M., Wratten, S. D., Landis, D. A., & You, M. (2017). Habitat management to suppress pest populations: Progress and prospects. *Annual Review of Entomology*, 62, 91–109. <https://doi.org/10.1146/annurev-ento-031616-035050>
- Gustianingtyas, M., Herlinda, S., Suwandi, Suparman, Hamidson, H., Hasbi, Setiawan, A., Verawaty, M., Elfita, & Arsi. (2020). Toxicity of entomopathogenic fungal culture filtrate of lowland and highland soil of South Sumatra (Indonesia) against *spodoptera litura* larvae. *Biodiversitas*, 21(5), 1839–1849. <https://doi.org/10.13057/biodiv/d210510>
- Hamid, H., Anjelina, S., Reffinaldon, R., Yunisman, Y., & Candra, E. (2025). The first report of the cassava lace bug (Hemiptera : Tingidae) and their impact on cassava plants on Sumatra Island , Indonesia. 0.
- Henuhili, V., & Aminatun, T. (2013). Konservasi Musuh Alami Sebagai Pengendali Hayati Hama Dengan Pengelolaan Ekosistem Sawah. *Jurnal Penelitian Saintek*, 18, 29–40.
- Herlina, G., Sidebang, M., H, J. R. B., Ayu, A., Pandansari, D., Dwiardana, A., Arya, I., Wibisana, P., Agosto, G., Adiel, Y., Sasongko, Y., Aditya, F., Saputra, R., Rachelia, N., & Ova, A. (2022). Pemanfaatan Hasil Alam Ubi Kayu Pada Desa Umbulrejo, Kecamatan Ponjong, Kabupaten Gunungkidul. 2(1).
- Herlinda, S., Efendi, R. A., Suharjo, R., Hasbi, Setiawan, A., Elfita, & Verawaty, M. (2020). New emerging entomopathogenic fungi isolated from soil in south Sumatra (Indonesia) and their filtrate and conidial insecticidal activity against *spodoptera litura*. *Biodiversitas*, 21(11), 5102–5113. <https://doi.org/10.13057/biodiv/d211115>
- Herlinda, S., Pujiastuti, Y., Irsan, C., Karenina, T., Budiarti, L., Rizkie, L., & Octavia, M. (2021). *Pengantar Ekologi Serangga*. Penerbit & Percetakan Universitas Sriwijaya (UNSRI).
- Hohbein, R. R., & Conway, C. J. (2018). Pitfall traps: A review of methods for estimating arthropod abundance. *Wildlife Society Bulletin*, 42(4), 597–606. <https://doi.org/10.1002/wsb.928>
- Hooper, D. U., Chapin III, F. S., Ewel, J. J., Hector, A., Inchausti, P., Lavorel, S., Lawton, J. H., Lodge, D. M., Loreau, M., Naeem, S., Schmid, B., Setälä, H., Symstad, A. J., Vandermeer, J., & Wardle, D. A. (2005). Effects of biodiversity on ecosystem functioning: A consensus of current knowledge.

Ecological Monographs, 75(1), 3–35. <https://doi.org/10.1890/04-0922>

- Joice, K. B. dos S., Tamara, T. dos S., Aleyres, B. C., Eliane, dos S., Diego, J. da S., Alverlan, da S. A., Rodrigo, A. P., Joao, P. F. B., Jhonatan, D. S. das N., Ana, P. G. da S. W., & Rubens, P. de B. (2019). Correlation of climatic elements with phases of the lace bug *Vatiga illudens* (Hemiptera: Tingidae) in two cassava cultivars (*Manihot esculenta* Crantz, Euphorbiaceae). *African Journal of Agricultural Research*, 14(10), 582–587. <https://doi.org/10.5897/ajar2018.13780>
- Kartohardjono, A. (2011). Komponen Pengendalian Hama Padi Berbasis Ekologi. *Jurnal Pengembangan Inovasi Pertanian*, 4(1), 29–46.
- Kiritani, K., & Kakiya, N. (1975). An analysis of the predator–prey system in the paddy field. *Researches on Population Ecology*, 17(1), 29–38. <https://doi.org/10.1007/BF02530737>
- Krebs, C. J. (2014). *Ecology: The Experimental Analysis of Distribution and Abundance* (6th ed.). Pearson Education Limited
- Landis, D. A., Wratten, S. D., & Gurr, G. M. (2000). Habitat management to conserve natural enemies of arthropod pests in agriculture. *Annual Review of Entomology*, 45, 175–201. <https://doi.org/10.1146/annurev.ento.45.1.175>
- Lawalata, J. J., & Anam, K. (2020). Pengamatan Jenis Predator Hama Tanaman Padi Di Kampung Karya Bumi Distrik Waibu Kabupaten Jayapura. *Median : Jurnal Ilmu Ilmu Eksakta*, 12(1), 13–20. <https://doi.org/10.33506/md.v12i1.824>
- Leong, J. M., & Thorp, R. W. (1999). Colour-coded sampling: The pan trap colour preferences of oligolectic and nonoligolectic bees associated with a vernal pool plant. *Ecological Entomology*, 24(3), 329–335. <https://doi.org/10.1046/j.1365-2311.1999.00196.x>
- Magurran, A. E. (2004). *Measuring Biological Diversity*. Blackwell Publishing
- Malado, M., Purnamasari, R., Nuryono, N., Monica, R. D., Lestari, S., Bahri, S., ... & Faizah, H. (2024). Pengendalian Hama dan Penyakit Tanaman Pertanian.
- Morris, E. K., Caruso, T., Buscot, F., Fischer, M., Hohberg, C., Sitting, D., ... & During, H. J. (2014). Choosing and using diversity indices: Insights for ecological applications from the German Biodiversity Exploratories. *Ecology and Evolution*, 4(18), 3514–3524. <https://doi.org/10.1002/ece3.1155>
- Nelly, N., Syahrawati, M. Y., dan Hamid, H. (2017). Abundance of corn planthopper (*Stenocranus pacificus*) (hemiptera: Delphacidae) and the potential natural enemies in West Sumatra, Indonesia. *Biodiversitas*, 18(2), 696–700. <https://doi.org/10.13057/biodiv/d180236>
- Neuenschwander, P., Hammond, W. N. O., Gutierrez, A. P., Cudjoe, A. R., Adjakloe, R., Baumgartner, J. U., & Regev, U. (1989). Impact assessment of the biological control of the cassava mealybug, *Phenacoccus manihoti* Matile-Ferrero (Hemiptera: Pseudococcidae) by the introduced parasitoid *Epidinocarsis lopezi* (De Santis) (Hymenoptera: Encyrtidae). *Bulletin of Entomological Research*, 79(4), 579–594.

<https://doi.org/10.1017/S0007485300018733>

- Nurkomar, I., Trisnawati, D. W., Azhar, A., & Saputra, J. A. (2023). Biodiversity of insects parasitoid and predator in cassava (*Manihot esculenta*) agroecosystem in Indonesia. *Proceedings of the Zoological Society*
- Nurmasari, F. (2015). Keanekaragaman Kutu Putih Dan Musuh Alami Pada Tanaman Singkong (*Manihot esculenta* Crantz.) Universitas Jember, 35.
- Nurdjanah, S., Susilawati, & Sabatini, M. R. (2007). Prediksi Kadar Pati Ubi Kayu (*Manihot Esculenta*) Pada Berbagai Umur Panen Menggunakan Penetrometer. *Jurnal Teknologi Dan Industri Hasil Pertanian*, 12(2), 65–73.
- Nyffeler, M., & Birkhofer, K. (2017). An estimated 400–800 million tons of prey are annually killed by the global spider community. *The Science of Nature*, 104(3–4), 30. <https://doi.org/10.1007/s00114-017-1440-1>
- Offenberg, J. (2015). Ants as tools in sustainable agriculture. *Journal of Applied Ecology*, 52(5), 1197–1205. <https://doi.org/10.1111/1365-2664.12496>
- Perfecto, I., & Vandermeer, J. (2020). The assembly and importance of a novel ecosystem: The ant community of coffee farms in Puerto Rico. *Ecology and Evolution*, 10(23), 12650–12662. <https://doi.org/10.1002/ece3.6785>
- Puspitarini, R. D., Fernando, I. T. O., Setiawan, Y., & Anggraini, D. (2021). Short Communication : First record of the cassava lace bug *Vatiga illudens* (Drake , 1922) (Hemiptera : Heteroptera : Tingidae) from East Java , Indonesia. 22(7), 2870–2876. <https://doi.org/10.13057/biodiv/d220738>
- Refiana, F., Triatmoko, E., & Fitriadi, S. (2021). Produktivitas Dan Pendapatan Usaha Tani Ubi Kayu (*Manihot Utilisima*) Di Desa Tungkaran Kabupaten Banjar. *Ziraa'Ah Majalah Ilmiah Pertanian*, 46(2), 185. <https://doi.org/10.31602/zmip.v46i2.4583>
- Restiani, R., Roslim, D. I., & Herman. (2014). Karakter Morfologi Ubi Kayu (*Manihot Esculenta* Crantz) Hijau Dari Kabupaten Pelalawan. *Jom Fmipa*, 1(2), 619–623.
- Rosumek, F. B., Giehr, J., griebeler, E. M., & (2022). The effects of ants on pest control: A meta-analysis. *Proceedings of the Royal Society B: Biological Sciences*, 289(1978), 20221316. <https://doi.org/10.1098/rspb.2022.1316>
- Rukmana, R. (1997). Ubi Kayu Budidaya dan Pasca panen. Kanisius. Yogyakarta
- Schulthess, F., Baumgärtner, J. U., Delucchi, V., & Gutierrez, A. P. (1991). The influence of the cassava mealybug, *Phenacoccus manihoti* Mat.-Ferr. (Horn., Pseudococcidae) on yield formation of cassava, *Manihot esculenta* Crantz. *Journal of Applied Entomology*, 111(1–5), 155–165. <https://doi.org/10.1111/j.1439-0418.1991.tb00306.x>
- Sembel, I. D. T. (2010). Pengendalian Hayati, Hama-hama serangga tropis dan gulma.
- Symondson, W. O. C., Sunderland, K. D., & Greenstone, M. H. (2002). Can generalist predators be effective biocontrol agents? *Annual Review of Entomology*, 47, 561–594.

<https://doi.org/10.1146/annurev.ento.47.091201.145240>

- Suroto, A., Oktaviani, E., & Khairunnisai, A. (2024). Eksplorasi dan deskripsi musuh alami potensial hama kutu kepik (*Vatiga Iludens* Drake 1922) (Hemiptera: Tingidae) dan serangga yang berasosiasi pada pertanaman singkong (*Manihot Esculenta*) di Kabupaten Banyumas, Provinsi Jawa Tengah. *Gontor AGROTECH Science journal*, 9(1), 11–24. <https://doi.org/10.21111/agrotech.v10i1.10841>
- Thamrin, M., Mardhiyah, A., & Samsul Efendi Marpaung. (2013). Analisis Usahatani Ubi Kayu (*Manihot Utilissima*). *Agrium*, 18(1), 57–64.
- Tscharntke, T., Klein, A. M., Kruess, A., Steffan-Dewenter, I., & Thies, C. (2005). Landscape perspectives on agricultural intensification and biodiversity – ecosystem service management. *Ecology Letters*, 8(8), 857–874. <https://doi.org/10.1111/j.1461-0248.2005.00782.x>
- Wahyuni, S. H., Kuswardani, R. A., & Noer, Z. (2025). Serangga Penyerbuk pada Tanaman Aren di Kabupaten Tapanuli Selatan.
- Wahyurini, E., dan Sugandini, D. (2021). Budidaya Dan Aneka Olahsan Singkong.
- Wardani, N. (2015). Kutu Putih Ubi Kayu, *Phenacoccus Manihoti* Matile-ferrero (Hemiptera: Pseudococcidae), Hama Invasif Baru Di Indonesia. Bogor: IPB.
- Wyckhuys, K. A. G., Hughes, A. C., Buber, C., Czepak, C., Graziosi, I., Gross, J., & Heimpel, G. E. (2018). Continental-scale suppression of an invasive pest by a host-specific parasitoid underlines both environmental and economic benefits of arthropod biological control. *PeerJ*, 6, e5796. <https://doi.org/10.7717/peerj.5796>
- Yuliawati. (2009). Pengelolaan tanaman dan organisme pengganggu tanaman (OPT) ubi kayu (*Manihot esculenta* Crantz.) di Kecamatan Ciemas, Sukabumi Dan Kecamatan Dramaga, Bogor.
- Zeddies, J., Schaab, R. P., Neuenschwander, P., dan Herren, H. R. (2001). Economics of biological control of cassava mealybug in Africa. *Agricultural Economics*, 24(2), 209–219.
- Zhang, G. R., Gu, D. X., & Liu, X. (2016). Transcriptome response to temperature stress in the wolf spider *Pardosa pseudoannulata* (Araneae: Lycosidae). *Ecology and Evolution*, 6(11), 3540–3554.
- Zhao, J. Z. (2001). Studies on the wolf spider *Pardosa pseudoannulata* as an important predator of rice pests in China. *Acta Arachnologica Sinica*, 10(2), 111–116