

DAFTAR PUSTAKA

- Aramaki-Hattori, N. *et al.* (2017) 'Relationship between Keloid Formation and YAP/TAZ Signaling', *Plast Reconstr Surg Glob Open*, 5(6), pp. 1–3. doi:10.1097/GOX.0000000000001357.
- Arima, J. *et al.* (2015) 'Hypertension: A systemic key to understanding local keloid severity', *Wound Repair Reg*, 23(2), pp. 213–221. doi:10.1111/wrr.12277.
- Bai, D., Ueno, L. and Vogt, P. (2009) 'Akt-mediated regulation of NFkB for the oncogenicity of PI3K and Akt', *Int J Cancer*, pp. 2863–2870.
- Bhattacharya, S., Ray, R.M. and Johnson, L.R. (2005) 'STAT3-mediated transcription of Bcl-2, Mcl-1 and C-IAP2 prevents apoptosis in polyamine-depleted cells', *Biochem J*, 392(2), pp. 335–344. doi:10.1042/BJ20050465.
- Boyce, D.E. *et al.* (2001) 'Inflammatory cell subpopulations in keloid scars', *Br J Plast Surg*, 54(6), pp. 511–516. doi:10.1054/bjps.2001.3638.
- Chai, J., Li, K. and Zhang, Y. (2025) 'Transcriptomic Analysis of Pressure Therapy on Hypertrophic Scars', *Eur res med J*, inpress, pp. 1–19.
- Chen, B., Li, H. and Xia, W. (2020) 'The role of Th1/Th2 cell chemokine expression in hypertrophic scar', *Int Wound J*, 17(1), pp. 197–205. doi:10.1111/iwj.13257.
- Chen, C.Y. *et al.* (2018) 'PTEN: Tumor Suppressor and Metabolic Regulator', *Front. Endocrinol.*, 9(JUL), pp. 1–12. doi:10.3389/fendo.2018.00338.
- Chen, Y. *et al.* (2019) 'Connection between T regulatory cell enrichment and collagen deposition in keloid', *Exp Cell Res*, 383(2), p. 111549.
- Chen, Y. *et al.* (2025) 'miR-3606-3p alleviates skin fibrosis by integratively suppressing the integrin / FAK , p-AKT / p-ERK , and TGF- b signaling cascades', *J Adv Res*, 75, pp. 271–290. doi:10.1016/j.jare.2024.11.027.
- Chow, J.Y.C. *et al.* (2008) 'TGF- β mediates PTEN suppression and cell motility through calcium-dependent PKC- α activation in pancreatic cancer cells', *Am J Physiol Gastrointest Liver Physiol*, 294(4), pp. 2–20. doi:doi:10.1152/ajpgi.00411.2007.
- Chun, Q. *et al.* (2016) 'Dynamic biological changes in fibroblasts during hypertrophic scar formation and regression', *Int Wound J*, 13(2), pp. 257–262. doi:10.1111/iwj.12283.
- Daele, U. Van *et al.* (2022) 'Mechanomodulation: Physical Treatment Modalities Employ Mechanotransduction to Improve Scarring', *Eur Burn J*, 3(2), pp. 241–255.
- Darby IA and Hewitson TD (2007) 'Fibroblast differentiation in wound healing and fibrosis', *Int Rev Cytol*, 257, pp. 143–79.
- Dirand, Z. *et al.* (2024) 'Macrophage phenotype is determinant for fibrosis development in keloid disease', *Matrix Biol*, 128(February), pp. 79–92.

doi:10.1016/j.matbio.2024.03.001.

Do, D. V. *et al.* (2012) 'Interleukin-18 system plays an important role in keloid pathogenesis via epithelial-mesenchymal interactions', *Br J Dermatol*, 166(6), pp. 1275–1288. doi:10.1111/j.1365-2133.2011.10721.x.

Dohi, T. *et al.* (2015) 'Tissue inhibitor of metalloproteinase-2 suppresses collagen synthesis in cultured keloid fibroblasts', *Plast Reconstr Surg Glob Open*, 3(9), pp. 1–11. doi:10.1097/GOX.0000000000000503.

Duan, X. *et al.* (2020) 'Identification and analysis of dysregulated lncRNA and associated ceRNA in the pathogenesis of keloid', *Ann Transl Med*, 3(1), pp. 1–11. doi:10.21037/atm.2020.01.07.

Eaton, R.G. and Sperling, R.A. (2007) *Grabb and Smith Plastic Surgery*. sixth. Edited by S.S. Charles Thorne, Robert Beasley, Sherrell Aston, Scott Bartlett, Geoffrey Gurtner. Philadelphia: Lippincott Williams & Wilkins.

De Felice, B. *et al.* (2007) 'Differential p63 and p53 expression in human keloid fibroblasts and hypertrophic scar fibroblasts', *DNA and Cell Bio*, 26(8), pp. 541–547. doi:10.1089/dna.2007.0591.

Feng, J. and Mertz, B. (2015) 'Novel Phosphatidylinositol 4 , 5-Bisphosphate Binding Sites on Focal Adhesion Kinase', *PLoS One*, 10(7), pp. 1–12. doi:10.1371/journal.pone.0132833.

Fratczak, A. *et al.* (2025) 'Botulinum toxin on a mission to erase keloids : does the dream come true ?', *Adv Dermatol Allergo*, 42(5), pp. 433–438.

Fujita, M. *et al.* (2019) 'NEDD4 Is Involved in Inflammation Development during Keloid Formation', *J Invest Dermatol*, 139(2), pp. 333–341. doi:10.1016/j.jid.2018.07.044.

Ghazawi, F.M. *et al.* (2018) 'Insights into the Pathophysiology of Hypertrophic Scars and Keloids: How Do They Differ?', *Adv Skin Wound Care*, 31(1), pp. 582–595. doi:doi: 10.1097/01.ASW.0000527576.27489.

Glaviano, A. *et al.* (2023) 'PI3K/AKT/mTOR signaling transduction pathway and targeted therapies in cancer', *Mol Cancer*, 22(1), pp. 1–37. doi:10.1186/s12943-023-01827-6.

Goyal, A. *et al.* (2016) 'Differential expression of Hedgehog and Snail in cutaneous fibrosing disorders implications for targeted inhibition', *Am J Clin Pathol*, 146(6), pp. 709–717. doi:10.1093/AJCP/AQW192.

Guo, L., Chen, L. and Bi, S. (2012) 'PTEN inhibits proliferation and functions of hypertrophic scar fibroblasts', *Mol Cell Biochem*, 361, pp. 161–168. doi:10.1007/s11010-011-1100-2.

He, T. *et al.* (2021) 'Phosphatase and Tensin Homolog in Non-neoplastic Digestive Disease: More Than Just Tumor Suppressor.', *Front physio*, 12, pp. 1–15. doi:10.3389/fphys.2021.684529.

Horibe, K., Hara, M. and Nakamura, H. (2021) 'M2-like macrophage infiltration and transforming growth factor- β secretion', *Arch Oral Biol*, 123(105042), pp. 1–8. doi:10.1016/j.archoralbio.2021.105042.

Horng, H. *et al.* (2017) 'Estrogen Effects on Wound Healing', *Int. J. Mol. Sci.*, 18, pp. 1–14. doi:10.3390/ijms18112325.

Hsu, C. *et al.* (2018) 'Caveolin-1 Controls Hyperresponsiveness to Mechanical Stimuli and Activation in Keloid Fibroblasts', *J Invest Dermatol*, 138, pp. 208–218. doi:10.1016/j.jid.2017.05.041.

Huang, C. and Ogawa, R. (2014) 'The link between hypertension and pathological scarring: Does hypertension cause or promote keloid and hypertrophic scar pathogenesis?', *Wound Repair Regen*, 22(4), pp. 462–466. doi:10.1111/wrr.12197.

Huang, Y. *et al.* (2023) 'Enhancement of Zyxin Promotes Skin Fibrosis by Regulating FAK / PI3K / AKT and TGF- β Signaling Pathways via Integrins', *Int J Biol Sci*, 19(8), pp. 2394–2408. doi:10.7150/ijbs.77649.

Ibrahim, N.E., Shaharan, S. and Dheansa, B. (2020) 'Adverse Effects of Pregnancy on Keloids and Hypertrophic Scars Case Presentation', *Cureus*, 12(12), pp. 1–9. doi:10.7759/cureus.12154.

Imaizumi, R. *et al.* (2009) 'Promoted activation of matrix metalloproteinase (MMP)-2 in keloid fibroblasts and increased expression of MMP-2 in collagen bundle regions: Implications for mechanisms of keloid progression', *Histopathology*, 54(6), pp. 722–730. doi:10.1111/j.1365-2559.2009.03287.x.

Jin, Q. *et al.* (2018) 'Macrophages in keloid are potent at promoting the differentiation and function of regulatory T cells', *Exp Cell Res*, 362(2), pp. 472–476. doi:10.1016/j.yexcr.2017.12.011.

Jumper, N. *et al.* (2017) 'A Role for Neuregulin-1 in Promoting Keloid Fibroblast Migration', *Acta Derm Venereol*, 97(6), pp. 675–684. doi:10.2340/00015555-2587.

Karin, M. (2009) 'NF- κ B as a Critical Link Between Inflammation and Cancer', *Cold Spring Harb Perspect Biol*, 1(5), pp. 1–11.

Katzung, B.G. (2018) *Basic & Clinical Pharmacology*. Fourteenth. Edited by B.G. Katzung. California: McGraw-Hill Education.

Kim, H. Do *et al.* (2013) 'Recurrent auricular keloids during pregnancy', *Arch Plastic Surg*, 40(1), pp. 70–72. doi:10.5999/aps.2013.40.1.70.

Kotian, S. *et al.* (2017) 'Dual Inhibition of HDAC and Tyrosine Kinase Signaling Pathways with CUDC-907 Inhibits Thyroid Cancer Growth and Metastases CUDC-907 Inhibits Thyroid Cancer Growth and Metastases', *Clin Cancer Res*, 23(17), pp. 5044–5054.

Kuehlmann, B. *et al.* (2020) 'Mechanotransduction in Wound Healing and Fibrosis', *J Clin Med*, 9(5), p. 1423. doi:10.3390/jcm9051423.

Kumar, V., Abbas, A.K., Fausto, N. and Mitchell, R.. (ed.) (2007) *Robbins Basic Pathology*. 8th editio. Philadelphia: Saunders Elsevier.

Lee, J.Y.Y. *et al.* (2004) 'Histopathological differential diagnosis of keloid and hypertrophic scar', *Am J Dermatopathol.*, 26(5), pp. 379–384. doi:10.1097/00000372-200410000-00006.

Lee, S.Y. *et al.* (2020) 'IL-17 Induced Stromal Cell-Derived Factor-1 and Profibrotic Factor in Keloid-Derived Skin Fibroblasts via the STAT3 Pathway', *Inflammation*, 43(2), pp. 664–672. doi:10.1007/s10753-019-01148-1.

Lei, R. *et al.* (2019) 'HIF-1 α promotes the keloid development through the activation of TGF- β /Smad and TLR4/MyD88/ NF- κ B pathways', *Cell Cycle*, 18(23), pp. 1–11. doi:10.1080/15384101.2019.1670508.

Li, B.S.H.U. *et al.* (2016) 'Role of mechanical strain-activated PI3K / Akt signaling pathway in pelvic organ prolapse', *Mol Med Rep*, 14, pp. 243–253. doi:10.3892/mmr.2016.5264.

Li, Jing *et al.* (2014) 'Risk of Keloid Associated with Polymorphic PTEN Haplotypes in the Chinese Han Population', *Wounds*, 26(1), pp. 21–27.

Li, Q. *et al.* (2021) 'Increased sensitivity to TNF- α promotes keloid fibroblast hyperproliferation by activating the NF- κ B, JNK and p38 MAPK pathways', *Exp Ther Med*. doi:10.3892/etm.2021.9933.

Lichtman, M.K., Otero-Vinas, M. and Falanga, V. (2016) 'Transforming growth factor beta (TGF- β) isoforms in wound healing and fibrosis', *Wound Repair and Regen*, 24(2), pp. 215–222. doi:10.1111/wrr.12398.

Lim, C.P. *et al.* (2006) 'Stat3 contributes to keloid pathogenesis via promoting collagen production, cell proliferation and migration', *Oncogene*, 25(39), pp. 5416–5425. doi:10.1038/sj.onc.1209531.

Limandjaja, G.C. *et al.* (2020) 'The Keloid Disorder: Heterogeneity, Histopathology, Mechanisms and Models', *Front Cell Dev Biol.*, 8(May), pp. 1–24. doi:10.3389/fcell.2020.00360.

Linjawi, S.A., Tork, S.E. and Shaibah, R.M. (2016) 'Genetic association of the COL1A1 gene promoter –1997 G/T (Rs1107946) and sp1 +1245 G/T (rs1800012) polymorphisms and keloid scars in a Jeddah population', *Turk J Med Sci*, 46(2), p. 414. doi:10.3906/sag-1412-41.

Liu, J. *et al.* (2018) 'Methylation of secreted frizzled-related protein 1 (SFRP1) promoter downregulates Wnt / β -catenin activity in keloids', *J Mol Histol*, 49(2), pp. 185–193. doi:10.1007/s10735-018-9758-3.

Liu, J. *et al.* (2025) 'The PI3K / AKT / mTOR pathway in scar remodeling and keloid formation: mechanisms and therapeutic perspectives', *Front. Pharmacol*, 16. doi:10.3389/fphar.2025.1678953.

Liu, S.L. *et al.* (2018) 'Quantitative Lipid Imaging Reveals a New Signaling Function of Phosphatidylinositol-3,4-Bisphosphate: Isoform- and Site-Specific Activation of Akt', *Molecular Cell*, 71(6), pp. 1092-1104.e5. doi:10.1016/j.molcel.2018.07.035.

Liu, X. *et al.* (2022) 'Single-Cell RNA-Sequencing Reveals Lineage-Specific Regulatory Changes of Fibroblasts and Vascular Endothelial Cells in Keloids', *J Invest Dermatol*, 142(1), pp. 124-135.e11. doi:10.1016/j.jid.2021.06.010.

Liu, Y. *et al.* (2012) 'MiRNA Expression Profiles in Keloid Tissue and Corresponding Normal Skin Tissue', *Aesth Plast Surg*, 36(1), pp. 193–201.

doi:10.1007/s00266-011-9773-1.

Liu, Y. *et al.* (2014) 'MicroRNA-21 affects proliferation and apoptosis by regulating expression of PTEN in human keloid fibroblasts', *Plast Reconstr Surg*, 134(4), pp. 561e-573e. doi:10.1097/PRS.0000000000000577.

Liu, Y. *et al.* (2023) 'Diagnosis and Treatment of Keloid: Method Summary and Effect Evaluation', *Clin Cosmet Investig Dermatol*, 16, pp. 3775–3783. doi:10.2147/CCID.S446018.

Livak, K.J. and Schmittgen, T.D. (2001) 'Analysis of Relative Gene Expression Data Using Real- Time Quantitative PCR and the 2^{-ΔΔC_T} Method', *METHODS*, 25(2001), pp. 402–408. doi:10.1006/meth.2001.1262.

Longley, D.B., Harkin, D.P. and Johnston, P.G. (2003) '5-Fluorouracil : Mechanism of Action and Clinical Strategies', *Cancer*, 3(May), pp. 330–338. doi:10.1038/nrc1074.

Lu, S. *et al.* (2020) 'PTEN Protects Against Angiotensin II-Induced Pathological Vascular Fibrosis and Remodeling', *Arterioscler Thromb Vasc BioCancer Cell*, 40(2), pp. 394–403. doi:10.1161/ATVBAHA.119.313757.PTEN.

Lu, Y.Y. *et al.* (2021) 'GLUT-1 enhances glycolysis, oxidative stress, and fibroblast proliferation in Keloid', *Life*, 11(6), pp. 1–18. doi:10.3390/life11060505.

Lv, D. *et al.* (2023) 'S-Nitrosylation-mediated coupling of DJ-1 with PTEN induces PI3K / AKT / mTOR pathway-dependent keloid formation', *Burns & Trauma*, 11, pp. 1–14. doi:https://doi.org/10.1093/burnst/tkad024.

Lv, W. *et al.* (2020) 'Epigenetic modification mechanisms involved in keloid: current status and prospect', *Clin Epigenetics*, 12(1), pp. 1–19. doi:10.1186/s13148-020-00981-8.

Mandal, K. (2020) 'Review of PIP2 in cellular signaling, functions and diseases', *Int J Mol Sci*, 21(21), pp. 1–20. doi:10.3390/ijms21218342.

Marks, P.A., Miller, T. and Richon, V.M. (2003) 'Histone deacetylases', *Current opinion in pharmacology*, 3(4), pp. 344–351.

Marongiu, M. *et al.* (2016) 'Novel action of FOXL2 as mediator of Colla2 gene autoregulation', *Dev Biol*, 416(1), pp. 200–211. doi:10.1016/j.ydbio.2016.05.022.

Marsh, D.J. *et al.* (1998) 'Mutation spectrum and genotype-phenotype analyses in Cowden disease and Bannayan – Zonana syndrome , two hamartoma syndromes with germline PTEN mutation', *Hum Mol Genet*, 7(3), pp. 507–515.

Messadi, D. V. *et al.* (2004) 'Activation of NFκB signal pathways in keloid fibroblasts', *Arch Dermatol Res*, 296(3), pp. 125–133. doi:10.1007/s00403-004-0487-y.

Mori, R. *et al.* (2014) 'Reduced FOXO1 Expression Accelerates Skin Wound Healing and Attenuates Scarring', *Am J Pathol*, 184(9), pp. 2465–2479. doi:10.1016/j.ajpath.2014.05.012.

Moshref S S, M.S.T. (2010) 'Keloid and Hypertrophic Scars: Comparative Histopathological and Immunohistochemical Study', *JKAU: Med. Sci*, 17(3), pp. 3–22. doi:10.4197/Med. 17-3.1.

Murakami, T. (2024) 'Pharmacotherapy for Keloids and Hypertrophic Scars', *Int J Mol Sci*, 25(9), pp. 1–61.

Murao, N. *et al.* (2014) 'Treg-enriched CD4⁺ T cells attenuate collagen synthesis in keloid fibroblasts', *Exp Dermatol*, 23(4), pp. 266–271. doi:10.1111/exd.12368.

Nakajima, Y. *et al.* (2022) 'Mast Cells Are Activated in the Giant Earlobe Keloids: A Case Series', *Int J Mol Sci*, 23(18), pp. 1–7. doi:10.3390/ijms231810410.

Nakashima, M. *et al.* (2010) 'A genome-wide association study identifies four susceptibility loci for keloid in the Japanese population', *Nat Genet*, 42(9), pp. 768–772. doi:10.1038/ng.645.

Nangole, F.W. *et al.* (2021) 'Multiple Cytokines Elevated in Patients with Keloids: Is It an Indication of Auto-Inflammatory Disease? Multiple Cytokines Elevated in Patients with Keloids: Is It an Indication of Auto-Inflammatory Disease?', *J Inflamm Res*, 14, pp. 2465–2470. doi:10.2147/JIR.S312091.

Nikolaos, G. and Frangogiannis (2020) 'Transforming growth factor- β in tissue fibrosis', *J Exp Med*, 217(3), pp. 1–36.

Noishiki, C., Hayasaka, Y. and Ogawa, R. (2019) 'Sex Differences in Keloidogenesis: An Analysis of 1659 Keloid Patients in Japan', *Dermatol Ther*, 9(4), pp. 747–754. doi:10.1007/s13555-019-00327-0.

Ogawa, R. *et al.* (2012) 'The relationship between skin stretching/contraction and pathologic scarring: The important role of mechanical forces in keloid generation', *Wound Repair Reg*, 20(2), pp. 149–157. doi:10.1111/j.1524-475X.2012.00766.x.

Ogawa, R. (2017) 'Keloid and Hypertrophic Scars Are the Result of Chronic Inflammation in the Reticular Dermis', *Int. J. Mol. Sci.*, 18(3), pp. 2–10. doi:10.3390/ijms18030606.

Ogawa, R., Akaishi, S. and Izumi, M. (2009) 'Histologic Analysis of Keloids and Hypertrophic Scars', *Ann Plast Surg*, 62(1), pp. 104–105. doi:10.1097/SAP.0b013e3181855172.

Ogawara, Y. *et al.* (2002) 'Akt enhances Mdm2-mediated ubiquitination and degradation of p53', *J Bio Chem*, 277(24), pp. 21843–21850. doi:10.1074/jbc.M109745200.

Patop, I.L., Wüst, S. and Kadener, S. (2019) 'Past, present, and future of circ RNA s', *EMBO J*, 38(16), p. e100836.

Peng, Y. *et al.* (2022) 'PI3K / Akt / mTOR Pathway and Its Role in Cancer Therapeutics: Are We Making Headway?', *Front Oncol*, 12, pp. 1–17. doi:10.3389/fonc.2022.819128.

Prabahar, A. *et al.* (2024) 'Unraveling the complex relationship between

mRNA and protein abundances : a machine learning-based approach for imputing protein levels from RNA-seq data', *NAR Genomics and Bioinformatics*, 6(February), pp. 1–12.

Qin, H. *et al.* (2022) 'EGR1/NOX4 pathway regulates oxidative stress and further facilitates fibrosis progression in keloids responses to TGF- β 1', *J Dermatol Sci*, 108(3), pp. 138–145. doi:10.1016/j.jdermsci.2022.12.009.

Richards, S. *et al.* (2015) 'ACMG Standards and Guidelines Standards and guidelines for the interpretation of sequence variants: a joint consensus recommendation of the American College of Medical Genetics and Genomics and the Association for Molecular Pathology', *Genetics in medicine*, 17(5), pp. 405–424. doi:10.1038/gim.2015.30.

Roth, S.Y., Denu, J.M. and Allis, C.D. (2001) 'Histone acetyltransferases', *Annu Rev Biochem*, 70(1), pp. 81–120.

Sang, P.F. *et al.* (2015) 'NEDD4-1 and PTEN Expression in Keloid Scarring', *Genetics and Molecular Research*, 14(4), pp. 13467–13475. doi:10.4238/2015.October.28.7.

Schmid, P. *et al.* (1998) 'Enhanced expression of transforming growth factor- β type I and type II receptors in wound granulation tissue and hypertrophic scar', *Am J Pathol*, 152(2), pp. 485–493.

Scully, M.M. *et al.* (2014) 'Rapid Estrogen Signaling Negatively Regulates PTEN Activity Through Phosphorylation in Endometrial Cancer Cells', *Hormones and Cancer*, 5(4), pp. 218–231. doi:10.1007/s12672-014-0184-z.

'SensiFAST™ Probe No - ROX Kit' (no date), 49(0). Available at: https://www.bioline.com/mwdownloads/download/link/id/2686/sensifast_probe_no_rox_kit_manual.pdf.

SensiFAST™ SYBR® No - ROX Kit (no date). Available at: https://www.bioline.com/mwdownloads/download/link/id/2754/sensifast_sybr_no_rox_kit_manual.pdf.

Seoudy, W.M. *et al.* (2023) 'Macrophages of the M1 and M2 types play a role in keloids pathogenesis', *Int Wound J*, 20(March 2022), pp. 38–45. doi:10.1111/iwj.13834.

Shaker, S.A., Ayuob, N.N. and Hajrah, N.H. (2011) 'Cell talk: A phenomenon observed in the keloid scar by immunohistochemical study', *Appl Immunohistochem Mol Morphol*, 19(2), pp. 153–159. doi:10.1097/PAI.0b013e3181efa2ef.

Shao, Y. *et al.* (2022a) 'Neutrophil extracellular traps contribute to myofibroblast differentiation and scar hyperplasia through the Toll-like receptor 9/nuclear factor Kappa-B/interleukin-6 pathway', *Burns & Trauma*, 10.

Shao, Y. *et al.* (2022b) 'Neutrophil extracellular traps contribute to myofibroblast differentiation and scar hyperplasia through the Toll-like receptor 9/nuclear factor Kappa-B/interleukin-6 pathway', *Burns and Trauma*, 10. doi:10.1093/burnst/tkac044.

Shi, J. *et al.* (2016) 'IL10 inhibits starvation-induced autophagy in

hypertrophic scar fibroblasts via cross talk between the IL10-IL10R-STAT3 and IL10-AKT-mTOR pathways', *Cell Death and Dis*, 7(3), pp. 1–11. doi:10.1038/cddis.2016.44.

Soares, E. and Zhou, H. (2018) 'Master regulatory role of p63 in epidermal development and disease', *Cell Mol Li Sci*, 75(7), pp. 1179–1190. doi:10.1007/s00018-017-2701-z.

Suh, B.-C. and Bertil Hille (2009) 'PIP2 is a necessary cofactor for ion channel function: How and why?', *Annu Rev Biophys*, 37, pp. 175–195. doi:10.1146/annurev.biophys.37.032807.125859.PIP.

Tang, Z. and Ding, J. (2020) 'MicroRNA-21 may be involved in the therapeutic effects of Galla chinensis ointment on keloid', *J Int Med Res* [Preprint]. doi:10.1177/0300060520909602.

Trace, A.P. *et al.* (2016) 'Keloids and Hypertrophic Scars : A Spectrum of Clinical Challenges', *Am J Clin Dermatol*, 17(3), pp. 201–223. doi:10.1007/s40257-016-0175-7.

Tsai, C.-H. and Ogawa, R. (2019) 'Keloid research: current status and future directions', *Scars Burn Heal.*, 5, pp. 1–8. doi:10.1177/2059513119868659.

Tu, T. *et al.* (2019) 'CUDC- 907 reverses pathological phenotype of keloid fibroblasts in vitro and in vivo via dual inhibition of PI3K/Akt/mTOR signaling and HDAC2', *Int J Mol Med*, 44(5), pp. 1789–1800. doi:DOI: 10.3892/ijmm.2019.4348.

Tuan, T.L. *et al.* (2003) 'Increased plasminogen activator inhibitor-1 in keloid fibroblasts may account for their elevated collagen accumulation in fibrin gel cultures', *Am J Pathol*, 162(5), pp. 1579–1589. doi:10.1016/S0002-9440(10)64292-7.

Vasudevan, K.M., Gurumurthy, S. and Rangnekar, V.M. (2004) 'Suppression of PTEN Expression by NF-kB Prevents Apoptosis', *Mol Cell Biol*, 24(3), pp. 1007–1021. doi:10.1128/MCB.24.3.1007.

Vogel, C. and Marcotte, E.M. (2013) 'Insights into the regulation of protein abundance from proteomic and transcriptomic analyses', *Nat Rev Genet*, 13(4), pp. 227–232. doi:10.1038/nrg3185.Insights.

Wang, Q., Wang, P., *et al.* (2021) 'Altered glucose metabolism and cell function in keloid fibroblasts under hypoxia', *Redox Biology*, 38(November 2020), p. 101815. doi:10.1016/j.redox.2020.101815.

Wang, Q., Xin, Y., *et al.* (2021) 'PI3K / AKT Pathway Promotes Keloid Fibroblasts Proliferation by Enhancing Glycolysis Under Hypoxia', *Wound Repair Regen.*, 31(2), pp. 1–14. doi:https://doi.org/10.21203/rs.3.rs-1022613/v1.

Wang, Z.C. *et al.* (2020) 'The Roles of Inflammation in Keloid and Hypertrophic Scars', *Front Immunol*, 11(December), pp. 1–10. doi:10.3389/fimmu.2020.603187.

Wen, W. *et al.* (2017) 'Inhibition of NEDD4 inhibits cell growth and invasion and induces cell apoptosis in bladder cancer cells', *Cell Cycle*, 16(16), pp. 1509–1514. doi:10.1080/15384101.2017.1338220.

Whitney, A.R. *et al.* (2003) 'Individuality and variation in gene expression patterns in human blood', *Proc Natl Acad Sci U S A*, 100(4), pp. 1896–1901. doi:doi10.1073pnas.252784499.

Wu, J. *et al.* (2020) 'RNA Sequencing Keloid Transcriptome Associates Keloids JAK3-Skewing', *Front Immunol*, 11(November), pp. 1–11. doi:10.3389/fimmu.2020.597741.

Xie, R. *et al.* (2024) 'Identification of potential therapeutic target SPP1 and related RNA regulatory pathway in keloid based on bioinformatics analysis', *Annals of Medicine*, 56(1), p. doi:10.1080/07853890.2024.2382949.

Xiong, S. (2013) 'Mouse models of Mdm2 and Mdm4 and their clinical implications', *Chinese Journal of Cancer*, 32(7), pp. 371–375. doi:10.5732/cjc.012.10286.

Xu, Z. *et al.* (2019) 'The Differential Expression of miRNAs and a Preliminary Study on the Mechanism of miR-194-3p in Keloids', *Biomed Res Int*, 2019, pp. 1–10. doi:10.1155/2019/8214923.

Yang, Y. *et al.* (2022) 'Androgen-related disorders and hormone therapy for patients with keloids', *Chinese Journal of Plastic and Reconstructive Surgery*, 4(1), pp. 44–48. doi:10.1016/j.cjprs.2022.02.003.

Yanping, W. *et al.* (2021) 'LncRNA TRHDE-AS1 inhibit the scar fibroblasts proliferation via miR-181a-5p/PTEN axis', *J Mol Histol*, 52(2), pp. 420–426. doi:https://doi.org/10.1007/s10735-021-09968-y.

Yazdani, Y. *et al.* (2016) 'The prognostic effect of PTEN expression status in colorectal cancer development and evaluation of factors affecting it: miR-21 and promoter methylation', *J Biomed Sci*, pp. 1–8. doi:10.1186/s12929-016-0228-5.

Yu, D. *et al.* (2016) 'Wnt.β-Catenin Signaling Exacerbates.pdf', *Cell Phys Biochem*, 39(5), pp. 2001–2013. doi:https://doi.org/10.1159/000447896.

Zhang, G. *et al.* (2017) 'DNA methylation of the CDC2L1 gene promoter region decreases the expression of the CDK11p58 protein and reduces apoptosis in keloid fibroblasts', *Arch Dermatol Res*, 310(2), pp. 107–115. doi:10.1007/s00403-017-1801-9.

Zhang, H.Y. *et al.* (2013) 'PTEN Mutation, Methylation and Expression in Breast Cancer Patients', *Oncology Letters*, 6(1), pp. 161–168. doi:10.3892/ol.2013.1331.

Zhang, J. *et al.* (2014) 'The regulation of TGF-β/SMAD signaling by protein deubiquitination', *Protein Cell*, 5(7), pp. 503–516. doi:DOI 10.1007/s13238-014-0058-8.

Zhang, W. *et al.* (2024) 'Commensal microbiome dysbiosis elicits interleukin-8 signaling to drive fibrotic skin disease', *PNAS Nexus*, 3(7), pp. 1–15. doi:10.1093/pnasnexus/pgae273.

Zhang, Y. *et al.* (2019) 'Knockdown of FOXM1 inhibits activation of keloid fibroblasts and extracellular matrix production via inhibition of TGF-β1/ Smad pathway', *Life Sci*, 232. doi:10.1016/j.lfs.2019.116637.

Zhong, L. *et al.* (2018) 'Identification and integrated analysis of microRNA expression profiles in keloid', *J Cosmet Dermatol*, 17, pp. 917–924. doi:10.1111/jocd.12706.

Zhou, X. *et al.* (2003) 'Germline PTEN Promoter Mutations and Deletions in Cowden / Bannayan-Riley-Ruvalcaba Syndrome Result in Aberrant PTEN Protein and Dysregulation of the Phosphoinositol-3-Kinase / Akt Pathway', *Am. J. Hum. Genet*, 72, pp. 404–411.

Zhu, H. *et al.* (2014) 'MicroRNA-21 Regulates hTERT via PTEN in Hypertrophic Scar Fibroblasts', *PLoS One*, 9(5), pp. 1–10. doi:10.1371/journal.pone.0097114.

