

DAFTAR PUSTAKA

1. Noiles K, Vender R. Are all seborrheic keratoses benign?: Review of the typical lesion and its variants. *J Cutan Med Surg.* 2008;12(5):203–10.
2. Hafner C, Vogt T. Seborrheic keratosis. *Journal Der Deutschen Dermatologischen Gesellschaft.* 2008;6(8):664–79.
3. Mawu FO. Tumor jinak kulit pada wajah. *e-CliniC.* 2016;4(1):641-4.
4. Yeatman JM, Kilkenny M, Marks R. The prevalence of seborrhoeic keratoses in an Australian population: Does exposure to sunlight play a part in their frequency?. *Br J Dermatol.* 1997;137:411–4.
5. Memon AA, Tomenson JA, Bothwell J, Friedmann PS. Prevalence of solar damage and actinic keratosis in a Merseyside population. *Br J Dermatol.* 2000;142(6):1154–9.
6. Gefilem GA, Suling PL, Kapantouw MG. Profil tumor jinak kulit di poliklinik kulit dan kelamin RSUD Prof. Dr. R. Kandou Manado periode 2009-2011. *J E-Clin.* 2013;1(1):1-9.
7. Hidajat D, Hapsari Y, Hendrawan IW. Karakteristik penyakit kulit pada geriatri di poliklinik kulit dan kelamin RSUD Provinsi Nusa Tenggara Barat periode 2012-2014. 2017;6(4):7–13.
8. Tejada C dos S, Mendoza-Sassi RA, Almeida Jr HL de, Figueiredo PN, Tejada VF dos S. Impact on the quality of life of dermatological patients in southern Brazil. *An Bras Dermatol.* 2011;86(6):1113–21.
9. Del Rosso J Q. Closer Look at Seborrheic Keratoses. *J Clin aesthetic dermatology* March. 2017;10(3):16-25.
10. Karadag AS, Parish LC. The status of the seborrheic keratosis. *Clin Dermatol.* 2017;36(2):275-7.
11. Cimpean I, Theate I, Vanhooteghem O. Seborrheic keratosis evolution into squamous cell carcinoma: A truly modified sun-related tumor?. *Dermatol Rep.* 2019;11(1):19–21.
12. Kwon OS, Hwang EJ, Bae JH, Park HE, Lee JC, Youn JI, et al. Seborrheic keratosis in the Korean males: Causative role of sunlight. *Photodermatol Photoimmunol Photomed.* 2003;19(2):73–80.

13. Thomas VD, Snaveley NR, Lee KK, Swanson NA. Benign epithelial tumors, hemartomas, and hyperplasias. In: Goldsmith LA, Katz SI, Gilchrest BA, Paller AS, Leffel DJ, Wolff K, editors. *Fitzpatrick's Dermatology in General Medicine*. 8th ed. New York: McGraw-Hill Companies, 2012. p.1319-23.
14. Kammeyer, A., Luiten, R.M., Oxidation events and skin aging, *Ageing Research Reviews* (2015), <http://dx.doi.org/10.1016/j.arr.2015.01.001>.
15. Fawzy MM, Abdel Aal MAM, Mohamed El-Shahed AR. Epidemiological Study of Seborrheic Keratosis among Egyptians in Great Cairo. *Egypt J Hosp Med*. 2018;72(4):4378–81.
16. Heidenreich B, Denisova E, Rachakonda S, Sanmartin O, Dereani T, Hosen I, et al. Genetic alterations in seborrheic keratoses. *Oncotarget*. 2017;8(22):36639–49.
17. Tsambaos D, Monastirli A, Kapranos N, Georgiou S, Pasmatzis E, Stratigos A, et al. Detection of human papillomavirus DNA in nongenital seborrheic keratoses. *Arch Dermatol Res*. 1995;287:612–5.
18. Gushi A, Kanekura T, Kanzaki T, Eizuru Y. Detection and sequences of human papillomavirus DNA in nongenital seborrheic keratosis of immunopotent individuals. 2003;1811:143–9.
19. Hunter J, Savin J, Dahl M, editors. Skin tumours. Seborrheic keratosis. In: *Clinical dermatology*. 3rd ed. Oxford: Blackwell Science, 2002. p. 255-6.
20. Roh NK, Hahn HJ, Lee YW, Choe YB, Ahn KJ. Clinical and Histopathological Investigation of Seborrheic Keratosis. *Ann Dermatol*. 2016;28(2):152–8.
21. Chellamaiah V, Vijayalakshmi B, Manoharan K. Clinico pathological study of cutaneous tumours of head & neck. *J Evol Med Dent Sci*. 2017;6(3):238–44.
22. James WD, Berger TG, Elston DM, editors. Epidermal nevi, neoplasm, and cysts. Seborrheic keratosis. In: *Andrew's diseases of the skin, clinical dermatology*. 11th ed. Elsevier Saunders, 2011. p. 625-7.
23. Nakamura S, Nishioka K. Enhanced expression of p16 in seborrheic keratosis: a lesion of accumulated senescent epidermal cells in G1 arrest.

- Br J Dermatol. 2003;149:560–5.
24. Atadja P, Wong H, Garkavtsev I, Veillette C, Riabowol K. Increased activity of p53 in senescing fibroblasts. *Proc Natl Acad Sci.* 1995;92:8348–52.
 25. Logie A, Dunois-larde C, Gasc J, Jorcano J, Werner S, Blanche M, et al. Activating mutations of the tyrosine kinase receptor FGFR3 are associated with benign skin tumors in mice and humans. *Human Molecular Genetics.* 2005;14(9):1153–60.
 26. Yuniati R, editors. Keratosis seboroik. In: *Buku ajar ilmu kesehatan kulit & kelamin pada layanan primer.* Universitas Diponegoro, 2017. p. 189-95.
 27. Hafner C, Oers JMMV, Hartmann A, Landthaler M, Stoehr R, Blaszyk H, et al. High frequency of FGFR3 mutations in adenoid seborrheic keratoses. *J Investigative Dermatol.* 2006;126:2404–7.
 28. Manaka I, Kadono S, Kawashima M, Kobayashi T, Imokawa G. The mechanism of hyperpigmentation in seborrhoeic keratosis involves the high expression of endothelin-converting enzyme-1 and TNF- α , which stimulate secretion of endothelin 1. *Brit J Dermatol.* 2001;145:895–903.
 29. Wolff K, Johnson RA, editors. Benign neoplasm and hyperplasias. Seborrheic keratosis. In: *Fitzpatrick's color atlas and synopsis of clinical dermatology.* 6th ed. New York: McGraw-Hill Companies, 2009. p. 215-8.
 30. Wardhana M. Dermoskopi: Cara non-invasif diagnostik lesi berpigmen. *Berkala Ilmu Kesehatan Kulit & Kelamin.* 2011;23(2):166–73.
 31. Braun RP, Rabinovitz H, Oliviero M, Kopf AW, Saurat JH. Dermoscopic diagnosis of seborrheic keratosis. *Clin Dermatol.* 2002;20(3):270–2.
 32. Minagawa A. Dermoscopy – pathology relationship in seborrheic keratosis. *J Dermatol.* 2017;44:518–24.
 33. Nadjar M, Anwar AI, Wahab S. Hubungan keratosis seboroik dengan penyakit kulit lain. *Medicinus.* 2015;28(2):38–42.
 34. Vincent KD, Abramovitz W. Cryosurgical treatment of benign skin conditions. Seborrheic keratosis. In: *Abramovits W, Graham G, Shai YH, Strumia R, editors. Dermatologic Cryosurgery.* London: Springer Nature, 2016. p. 589-93.

35. Wollina U. Seborrheic keratoses – the most common benign skin tumor of humans: Clinical presentation and an update on pathogenesis and treatment options. *Open Access Maced J Med Sci.* 2018;6(11):2270–5.
36. Tanojo H, Yenny SW, Lestari S. Perbandingan terapi keratosis seboroik wajah dengan teknik split-face antara laser karbon dioksida dengan elektrodeseikasi. 2013;40(2):112–5.
37. Fitzpatrick RE, Goldman MP, Esparza JR. Laser treatment of benign pigmented epidermal lesions using a 300 nsecond pulse and 510 nm wavelength. *J Dermatol Surg Oncol.* 1993;19:341-7.
38. Chun EY, Lee JB, Lee KH. Focal Trichloroacetic acid peel method for benign pigmented lesions in dark-skinned patients. *Dermatol Surg.* 2004;30(4):512–6.
39. Herron MD, Bowen AR, Krueger GG. Clinical trial seborrheic keratoses: A study comparing the standard cryosurgery with topical calcipotriene , topical tazarotene , and topical imiquimod. *Inter J Dermatol.* 2004;43:300–2.
40. Mitsuhashi Y, Kawaguchi M, Hozumi Y, Kondo S. Topical vitamin D3 is effective in treating senile warts possibly by inducing apoptosis. *J Dermatol.* 2005;32(6):420–3.
41. Keaney TC. Aging in the Male Face : Intrinsic and Extrinsic Factors. *American Soc for Dermatol Surg.* 2016;1–7.
42. Stauffer BL, Westby CM, Desouza CA. Endothelin-1 , aging and hypertension. *Wolters Kluwer Health.* 2008;23:350-5.
43. Hafner C, Lo E, Luis NM, Ferna A, Ribe A, Mentzel T, et al. Oncogenic PIK3CA mutations occur in epidermal nevi and seborrheic keratoses with a characteristic mutation pattern. *PNAS J.* 2007;104(33):13450-4.
44. Modenese A, Korpinen L. Solar Radiation Exposure and Outdoor Work : An Underestimated Occupational Risk. *Int J Environ Res Pub Health.* 2018;15(2063):1–24.
45. McGregor JM, Rustin MHA. Human papillomavirus and skin cancer. *Postgrad Med J.* 1994;70(828):682–5.
46. Rajesh G, Thappa DM, Jaisankar TJ, Chandrashekar L. Spectrum of

- seborrhic keratoses in south Indians : A clinical and dermoscopic study. *Indian J Dermatol.* 2011;77(4):483-8.
47. Farage MA, Miller KW, Elsner P, Maibach HI. Structural characteristic of the aging skin. *Cutan and Ocular Toxic.* 2007;(4):343–57.
48. Okazaki M, Yoshimura K, Uchida G, Harii K. Correlation between age and the secretions of melanocyte stimulating cytokines in cultured keratinocytes and fibroblasts. *Brit J Dermatol.* 2005;1:23–9.
49. Hafner C, Hartmann A, Oers JMM Van, Stoehr R, Zwarthoff EC, Hofstaedter F, et al. FGFR3 mutations in seborrhic keratoses are already present in flat lesions and associated with age and localization. *Modern Pathology.* 2007;20:895–903.
50. Gill D, Dorevitch A, Marks R. The prevalence of seborrhic keratoses in people aged 15 to 30 years. *Arch Dermatol.* 2016;136:1–4.
51. Alapatt G, Sukumar D, Bhat M. A clinicopathological and dermoscopic correlation of seborrhic keratosis. *Indian J Dermatol.* 2016;61(6):622-7.
52. Rongioletti F, Corbella L, Rebora A. Multiple familial seborrhic keratoses. *Dermatol.* 1988;176:43-5.
53. Shariff VNSA, Balamurugan L, Saravanan N. Seborrhic keratosis with autosomal dominant inheritance. *Int J Res Med Science.* 2017;5(3):1138–41.
54. Kennedy C, Bajdik CD, Willemze R, Gruijl FR De, Bavinck JNB. The influence of painful sunburns and lifetime sun exposure on the risk of actinic keratoses, seborrhic warts, melanocytic nevi, atypical nevi, and skin cancer. *J Investigative Dermatol.* 2003;120(6):1087–93.
55. Imokawa G, Miyagishi M, Yada Y. Endothelin 1 as new melanogen. *J Investigative Dermatol.* 1995;105(1):32-7.
56. Engel A, Johanson ML, Haynes SG. Health effects of sunlight in the united states. *Arch Dermatol.* 2015;124:72-9.