

DAFTAR PUSTAKA

- Birdsall M, Ledger W, Groome N. Inhibin A and Activin A in the first Trimester of Human Pregnancy. *J clin Endocrinol Metab* 2008; 82:1557-60
- Blumenstein M, Mitchell MD, Groome NP, Keelan JA. Hypoxia inhibit activin A production by term villous trophoblast in vitro. *Plasenta* 2009 : 23 : 735-41
- Cunningham FG et al. (2014). Hypertensive Disorder in Pregnancy. Dalam C. F. al, *William Obstetrics 23rd Ed.* New York: McGraw-Hill Companies Inc.
- Cunningham, F. G., Leveno, K. J., Bloom, S. L., Hauth, J. C., Gilstrap, L., & Wenstrom, K. D. (2014). Pregnancy Hypertension. Dalam F. G. Cunningham, K. J. Leveno, S. L. Bloom, J. C. Hauth, L. Gilstrap, & K. D. Wenstrom (Penyunt.), *Williams Obstetrics (24th Edition ed.)*. New York: The McGraw-Hill Companies.
- Davidson E, Riley S, Robert S, Shearing C, Groome N. Maternal serum activin, inhibin, human chorionic gonadotropin and α -fetoprotein as second trimester predictors of preeclampsia. *BJOG* 2005; 110: 46-52
- Dekker G. Hypertension. Dalam: *High Risk Pregnancy 4th Edition*. Elsevier Saunders, Philadelphia. 2011.
- Eleazar Soto, e. a. (2012). Late-onset preeclampsia is associated with an imbalance of angiogenic and anti-angiogenic factors in patients with and without placental lesions consistent with maternal underperfusion. *The Journal of maternal fetal and neonatal medicine* , 498–507.
- Eruo FU, S. B. (2007). Hipertensive Diseases in Pregnancy. in H. J. Reece EA, *Clinical Obstetrics The Fetus and Mother 3rd Edition* (683-699). Massachusetts, USA: Blackwell Publising Ltd.
- Hamad, R. (2011). Cardiovascular Function and Biomarkers in Women with Preeclampsia. *Depart. Of Women's amd Children's health, Kalolinska Institute, Stockholm.*
- Gallinelli, Gennazeni AD, Matteo ML, Caruso A, Woodruff. Episodic secretion of activin A in pregnant women. *Euro J Endocrinol* 1996;135:340-4
- Hanisch CG, Schlebusch H, Schmolling J. Adhesion molecules, activin and inhibin candidates for the biochemical prediction of hypertensive disease in pregnancy. *Arch Gynecol Obstet* 2009 ; 270 : 110-15
- Jeffrey A, Tailor R, Schellenberg JC. Serum activin A, inhibin A, and follistatin concentrations in preeclampsia or small for gestational age pregnancies. *Elsevier* 2002; 99: 267-74

Jonathan Lai. Activin A in the second Trimester of Human Pregnancy. *Journal J clin Endocrinol Metab* 2013; 84:1767-70

Keelan JA, Tailor R, Schellenberg J, Groome N, Mitchel M, North R. Serum Activin A, Inhibin A and Follistatin Concentrations in Preeclampsia or Small for Gestational Age Pregnancies. *Am J Obstet Gynecol* 2007;99: 267-74

Ling Yu, Li Dong, Liao Q P, high levels of activin A detected in preeclamptic placenta induce trophoblast cell apoptosis by promoting nodal signaling. *J Clin Endocrinol Metab* 2012, 97(8): E1370-E1379

Lubis DN. Gambaran Kematian Ibu di Bagian Obstetri Ginekologi RSUP Dr. M. Djamil Padang Tahun 2011-2012. *Bagian Obstetri dan Ginekologi FK UNAND. Padang*: 2013.

Madazli R, Kuseyrioglu B, Uzun H, Uludag S. Prediction of preeclampsia with maternal mid trimester plasental growth factor, activin A, fibronectin and uterine artery doppler velocimetry. *Intel J. Gynecol Obstet* 2007 : 89 : 251-7

Madi J and Sulin D. (2003). Angka Kematian Pasien Preeklampsia dan Eklampsia di RS. Dr. M. Djamil Padang tahun 1998-2002. *Bagian Obstetri dan Ginekologi FK. Unand/RS. Dr. Djamil Padang* .

Meads Ca et all, Methods of prediction and prevention of pre-eclampsia: systematic reviews of accuracy and effectiveness literature with economic modeling, HTA, United Kingdom, 2008.

Muttukrishna S, Groome N, Redman, Sargent. Serum inhibin A and activin A are elevated prior to the onset of preeclampsia. *Hum Reprod* 2006; 15:1640-5.

Muttukrishna S, Knight PG, Groome N, Redman. Activin A and inhibin A as possible endocrine markers for pre-eclampsia. *The lancet* 1997. 349:1285-87

Pipkin FB. Maternal physiology. In: Edmonds DK, ed. *Dewhurst's Textbook of Obstetrics & Gynaecology*. Seventh edition ed. Massachusetts: Blackwell 2007:10-8.

Redman. Pregnancy-related hypertension. Dalam: *Maternal-Fetal Medicine Principles and Practice*. Edisi 5. Saunders. Philadelphia: 2012. 859-892.

Roshadi, R. (2004). Hipertensi Dalam Kehamilan. Dalam R. Hariadi, *Ilmu Kedokteran Fetomaternal* (hal. 494-500). Surabaya: Himpunan Kedokteran Fetomaternal Perhimpunan Obstetri dan Ginekologi Indonesia.

Silver HM, et al. Mechanism of increased maternal serum total activin A and inhibin A in preeclampsia. *J Soc Gynecol Investig*. 2008; 9: 308-12

Smith MC, Davison JM. The Kidney and Preeclampsia. In : Fiona Lyall and Michael Belfort. Preeclampsia Etiology and Clinical Practice. CAMBRIDGE University Press, 2007. New York. 339-356.

Stefano L, Florio P, Reis FM, Petraglia F. Expression and secretion of activin A: possible physiological and clinical implication. Euro J Endocrinol, 2001:225-36

valenzuela F.J, e. a. (2012). Pathogenesis of Preeclampsia : The Genetic Component. *Journal of Pregnancy* .

Wallace E, Kolsky M, Edwards, Baker, Jenkin G. Maternal serum activin A levels in association with intrauterine fetal growth restriction. BJOG 2009;110:306-10

Yang Liu et al. High Levels of Activin A Detected in Preeclamptic Placenta Induce Trophoblast Cell Apoptosis by Promoting Nodal Signaling. J Clin Endocrinol Metab 2012; 97(8):E1370–E1379

Zilfira D. Adiponektin pada Preeklamsia. Bagian Obstetri dan Ginekologi FK Unand/RS Dr. M. Djamil Padang, 2012.

