

## DAFTAR PUSTAKA

1. Peric S, Branimir C, Porin P, Ivan S, Hrvoje T. Automated Static Perimetry: The Influence of Myopic Anisometropia on Evaluation of Visual Field. *Coll Antropol.* 2012;37(1):75-81.
2. Curtin BJ, Whitmore WG. The Optics of Myopia. In: Duane's Foundation of Clinical Ophthalmology. Northwestern University: Philadelphia Harper & Row. 2009. pp 1-11.
3. Matamoros E, Ingrand P, Pelen F, Bentaleb Y, Weber M, et al. Prevalence of Myopia in France. *Med (United States).* 2015;94(45):1-6.
4. Saw SM, Gus G, David K, Mohamed F, Daniel W, et al. Prevalence Rates of Refractive Errors in Sumatera. *IOVS.* 2002;43(10):3174-80.
5. Sawada Y, Makoto A, Hitomi S, Makoto I, Toyota I, et al. Optic Disc Margin Anatomic Features in Myopic Eyes with Glaucoma with Spectral-Domain OCT. *AAO Journal.* 2018;125(12):1886-96.
6. Ohno-matsui K, Spaide RIF. Myopic Optic Neuropathy. In: Pathologic Myopia. New York: Springer; 2014. pp 293–309.
7. Park HYL, Kim YC, Jung Y, Park CK. Vertical Disc Tilt and Features of The Optic Nerve Head Anatomy are Related to Visual Field Defect in Myopic Eyes. *Sci Rep.* 2019;9(1):1–9.
8. Hao L, Hui X, Xue G, Xiaoyu X, Xing L. Measurement of Structural Parameters of the Lamina Cribosa in Primary Open Angle Glaucoma and Chronic Primary Angle Closure Glaucoma by Optical Coherence Tomography and Its Correlations with Ocular Parameters. *Ophthalmic Res.* 2019;22(8):1-10.
9. Kim YW, Jin J, Dai WK, Michael JAG, Jean MM, et al. Clinical Assessment of Lamina Cribosa Curvature in Eyes with Primary Open Angle Glaucoma. *PLOS ONE.* 2016;10(3):1-15.
10. Sigal IA, Bo W, Nicholas GS, Tadamichi A, et al. Recent Advances in OCT Imaging of The Lamina Cribosa. *Br J Ophthalmol.* 2014;4(98):134-9.
11. Kim DW, Jin WJ, Yong WK, Michael JAG, Jean MM, et al. Prelamina and Lamina Cribosa in Glaucoma Patients with Unilateral Visual Field Loss. *IOVS.* 2016;57(4):1662-70.
12. Garg P, Laxmi S, Rubie M, Mona L. A Study on Systemic Risk Factors for Primary Open Angle Glaucoma. *Pharmaceutical science.* 2014;4(2): 1-8.
13. Park SC, Brumm J, Furlanetto RL, Netto C, Liu Y, Tello C, et al. Lamina Cribosa Depth in Different Stages of Glaucoma. *Investig Ophthalmol Vis Sci.* 2015;56(3):2059–64.

14. Fayed AAE. Evaluation of The Lamina Cribrosa Morphology in Correlation to Retinal Nerve Fiber Layer Thickness. *Delta J Ophthalmol.* 2017;18(1):166–9.
15. Downs JC, Girkin CA. Lamina Cribrosa in Glaucoma. *Curr Opin Ophthalmol.* 2018;28(2):113–9.
16. Oh B, Lee EJ, Kim H, Girard MJA, Mari JM. Anterior Lamina Cribrosa Surface Depth in Open-Angle Glaucoma : Relationship with the Position of the Central Retinal Vessel Trunk. *PLOS One.* 2016;11(6):1–13.
17. Shoji T, Hiroto K, Masayuki S, Motoyushi B, Masanori H, et al. Correlation Between Lamina Cribrosa Tilt Angles, Myopia, and Glaucoma Using OCT with a Wide Bandwidth Femtosecond Mode-Locked Laser. *PLOS One.* 2014;9(12):1-13.
18. Turgut B. Lamina Cribrosa Measurements for Glaucomatous Eyes: A New Focus Point in Optical Coherence Tomography. *Adv OphthalmolVis Syst.* 2018;8(1):70-72.
19. Wang X, Kong X, Jiang C, Li M, Yu J. Is the peripapillary retinal perfusion related to myopia in healthy eyes ? A prospective comparative study. *BMJ Open.* 2016;6(1):1–7.
20. Ha A, Tai JK, Michael JA, Girard, Jean MM, et al. Baseline Lamina Cribrosa Curvature and Subsequent Visual Field Progression Rate in Primary Open Angle Glaucoma. *Ophthalmology.* 2018;125(18):1898-1906.
21. Kim M, Karine DB, Mark AS, Leona D, Philip PC. Lamina Depth and Thickness Correlate with Glaucoma Severity. *Indian Journal of Ophthalmology.* 2016;64(2):358-63.
22. Tiffani DR, Hendriati, Fitratul I. Perbedaan *Lamina Cribrosa Depth* dengan *Retinal Nerve Fiber Layer Thickness* Pada Mahasiswa Kedokteran dengan Miopia. *Universitas Andalas.* 2019.
23. Lopes FS, Igor M, Izabela A, Syril KD, Roberto MV, et al. Structure Function Relationship in Glaucoma Using Enhanced Depth Imaging Optical Coherence Tomography-Derived Parameters: A Cross-Sectional Observational Study. *BMC Ophthalmology.* 2019;19(52):1-8.
24. Sun J, Zhou X, Kang LY, Sun X, Sui H, et al. Prevalence and Risk Factors for Primary Open Angle Glaucoma in a Rural Northeast China Population: A Population-based Survey in Bin County, Harbin. *Eye.* 2012;26(12):283-91.
25. Xiao H, Xiao YV, Yi-MZ, Xing L. Age Related Changes of The Central Lamina Cribrosa Thickness, Depth and Prelaminar Tissue in Healthy Chinese Subjects. *Int J Ophthalmol.* 2018;11(11):1842-47.
26. Park HYL, So HJ, Chan KP et al. Enhanced Depth Imaging Detects Lamina Cribrosa Thickness Differences in Normal Tension Glaucoma and Primary Open Angle Glaucoma. *American Academy of Ophthalmology.* 2012;119(2):10-20.

27. Seo JH, Tae WK, Robert NW. Lamina Cribrosa Depth in Healthy Eyes. *Investig Ophthalmol Vis Sci.* 2014;55(3):1241-50.
28. Kwun Y, Gyule H, Yoon JC, Jong CH, Changwon K. Optic Disc Characteristics and Visual Field Progression in Normal Tension Glaucoma Patients with Tilted Optic Discs. *J Glaucoma.* 2016;25(11):901-7.
29. Chang R, Kuldev S. Detecting and Treating Glaucoma in a Myopic Patient Without High IOP. *Glaucoma Today.* 2016;5(1):12-14.
30. Ren R, Hongli Y, Stuart K, Gardiner, Brad F, et al. Anterior Lamina Cribrosa Surface Depth, Age, And Visual Field Sensitivity in The Portland Progression Project. *Investig Ophthalmol Vis Sci.* 2014;55(3):1531-39.
31. Kyari F, Mohammed MA, Richard W, Jennifer RE, Winifred N, et al. Risk Factor for Open Angle Glaucoma in Nigeria: Results from the Nigeria National Blindness and Visual Impairment Survey. 2016;16(78):1-12.
32. Takusagawa HL, Ambika H, Anna K, Kouros NM, Sanita R, et al. Swept Source OCT for Evaluating the Lamina Cribrosa. *Ophthalmology.* 2019;44(3):1-9.
33. Kim YN, Joong WS, Kyung RS. Relationship between Progressive Changes in Lamina Cribrosa Depth and Deterioration of Visual Field Loss in Glaucomatous Eyes. *Korean J Ophthalmol.* 2018;32(6):470-77.
34. Nitta K, Kazuhisa S, Ryotaro W, Gaku T. Is High Myopia a Risk Factor for Visual Field Progression or Disk Hemorrhage in Primary Open Angle Glaucoma?. *Clinical Ophthalmology.* 2017;11(2):599-604.
35. Aung T, Paul JF, Steve KS, Siw PC, Wee KL, et al. Automated Static Perimetry: The Influence of Myopia and Its Method of Correction. *Ophthalmology.* 2001;108(8):290-5.
36. Shoeibi N, Nasrin MS, Ramin D, Asieh E. Visual Field Assessment in High Myopia with and without Tilted Optic Disc. *Clinical and Experimental Optometry.* 2016;3(11):1-5.
37. Ohno-matsui K, Noriaki S, Kenjiro Y, Kengo H, Takeshi Y, et al. Long-term Development of Significant Visual Field Defect in Highly Myopic Eyes. *Am J Ophthalmol.* 2011;152(2):256-65.
38. Kuzhuppilly N, Shilpa P, Shibi D, Aditya D. Reability of Visual Field Index in Staging Glaucomatous Visual Field Damage. *Journal of Clinical and Diagnostic Research.* 2018;12(6):5-9.
39. Susanna R, Roberto MV. Staging glaucoma patient: why and how? *The Open Ophthalmology Journal.* 2009; 3(5):59-64.
40. Park HY, Lopilly P, Kyung EH, Chan KP. Impact of Age and Myopia on the Rate of Visual Field Progression in Glaucoma Patients. *Medicine.* 2016;95(21):1-7.
41. Lee EJ, Kim TW, Weinreb RN, Park KH, Kim SH, et al. Visualization of

- the Lamina Cribrosa Using Enhanced Depth Imaging Spectral-Domain Optical Coherence Tomography. *AJOPHT*. 2011;152(1):87-95.
42. Doshi A, Ken O, Kreidl, Lorinna L, Douglas KS. Nonprogressive Glaucomatous Cupping and Visual Field Abnormalities in Young Chinese Males. *Ophthalmology*. 2007;114(3):472-79.
  43. Daguman IJ, Delfin MS. Correction of Lamina Cribosa and Standard Automated Perimetry Finding in Glaucoma and Non Glaucoma Patients. *J Ophthalmic Stud*. 2018;2(1):1-5.
  44. Chihara E, Xiaou L, Jin D, Yasuyuki T, Masayuki A, Masanori H, et al. Severity Myopia as A risk Factor for Progressive Visual Field Loss in Primary Open Angle Glaucoma. *Ophthalmologica*. 1997;211(2):66-71.
  45. Benjamin WJ. Refractive Status of the Eye. In: Borish Clinical Refraction. Missouri: Elsevier Inc; 2006. pp. 4–9.
  46. Cantor LB, Rapuano CJ, Cioffi GA. Optic of the Human Eye. In: Clinical optics. San Fransisco: American Academy of Ophthalmology; 2019-2020. pp. 161-181.
  47. Holden BA, Fricke TR, Wilson DA, Jong M, Naidoo KS, Sankaridurg P, et al. Global Prevalence of Myopia and High Myopia and Temporal Trends from 2000 through 2050. *Ophthalmology*. 2016;123(5):1036–42.
  48. Cantor LB, Rapuano CJ, Cioffi GA. Clinical Evaluation. In: Glaucoma. San Fransisco: American Academy of Ophthalmology; 2019-2020. pp. 57-66.
  49. Rand AR. The Basic Aspect of Glaucoma. In: Shields Textbook of Glaucoma. North Carolina: Lippincott Williams & Wilkins; 2011. pp. 83-90.
  50. Cantor LB, Rapuano CJ, Cioffi GA. Diagnostic Approach to Retinal Disease. In: Retina and Vitreus. San Fransisco: American Academy of Ophthalmology; 2019-2020. pp. 51-55.
  51. Barton J, Benatar M. Automated Perimetry (Humphrey Field Analyzer). In: Field of Vision: A Manual and Atlas of Perimetry. New Jersey: Humana Press; 2003. pp 45-56.
  52. Cantor LB, Rapuano CJ, Cioffi GA. Clinical Evaluation. In: Glaucoma. San Fransisco: American Academy of Ophthalmology; 2019-2020. pp. 78-99.
  53. Wild J. Short Wavelength Automated Perimetry. *Acta Ophthalmol Scand*. 2001;79:546–59.
  54. Gupta A, Choudhry R, Tandon C. Humphrey Field Analyzer. In: Step by Step Visual Field Examination. New Delhi: Jaypee Brothers; 2007. pp 133-56.
  55. Kahook BYMY, Noecker RJ. How Do You Interpret a 24-2 Humphrey Visual Field Printout? *Glaucoma Today*. 2007;12(2):57–9.
  56. Heijl A, Patella V, Bengtsson B. STATPAC Analysis of Single Field. In:

- The Field Analyzer Primer: Effective Perimetry. California: Carl Zeiss Meditec; 2012. pp 45–59.
57. Sankar PS, Laura O, Joan OB. The SCHEIE Visual Field Grading System. *Journal of clinical & experimental ophthalmology*. 2017;8(3):1-16.
  58. Handerer JD. Interpreting Visual Fields. *Glaucoma Today*. 2004;7(1):27-32.
  59. Day M, Duffy LA. Myopia and defocus: The current understanding. *Scand J Optom Vis Sci*. 2011;4(1):1-14.
  60. Cooper J, Tkatchenko A V. A Review of Current Concepts of the Etiology and Treatment of Myopia. *Eye Contact Lens*. 2018;44(4):231–47.
  61. Sood RS. Influence of Gender on The Prevalence of Myopia in Young Adults. *International Journal of Basic and Applied Medical Sciences*. 2012;2(1):201-204.
  62. Inoue R, Hangai M, Kotera Y, Nakanishi H, Mori S, Morishita S, et al. Coherence Tomography Imaging of Lamina Cribrosa in Glaucoma. *OPHTHA*. 2009;116(2):214–22.
  63. Srinivasan VJ, Dimitrova N, Zamudio JR, Jong RM, Soukup D, Resnick R, et al. Ultrahigh-Speed Optical Coherence Tomography for Three-Dimensional and En Face Imaging of the Retina and Optic Nerve Head. *PLoS One*. 2017;32(7):736–40.
  64. Jonas JB, Kutscher JN, Panda JS, Hayreh SS. Lamina Cribrosa Thickness Correlated with Posterior Scleral Thickness and Axial Length in Monkeys. *Acta Ophthalmol*. 2016;27(9):693-6.
  65. Jonas JB, Xu L. Histological Changes of High Axial Myopia. *Eye*. 2014;28(10):113-7.
  66. Jonas JB, Berenshtein E, Holbach L. Lamina Cribrosa Thickness and Spatial Relationships between Intraocular Space and Cerebrospinal Fluid Space in Highly Myopic Eyes. *Invest Ophthalmol Vis Sci*. 2004;45(8):2660-5.
  67. Yun S, Hahn IK, Sung KR, Yoon JY. Lamina Cribrosa Depth According to The Level of Axial Length in Normal and Glaucomatous Eyes. *Graefe's Arch Clin Exp Ophthalmol*. 2015;27(8):1-7.
  68. Juyal A, Punam V, Sanjeev M. A Comparatve Study of Visual Field in Emmetropic and Mupoic Subjects by Automated Perimetry. *Indian Journal of Applied Research*. 2017;7(5):611-3.
  69. Yuan ZY, Chen LF, Bao YL, Min QS, Xiu PC. The Relationship between Visual Field Global Indices and Retinal Nerve Fiber Layer Thickness in Healthy Myopes. *Journal of Ophthalmology*. 2014;10(11):1-8.
  70. Sharif NM, Nasser S, Asieh E, David A. Structure Versus Function in High Myopia Using Optical Coherence Tomography and Automated

- Perimetry. *Clinical and Experimental Optometry*. 2019;102(13):335-40.
71. Sawada Y, Masanori H, Makoto I, Takeshi Y. Association of Myopic Optic Disc Deformation with Visual Field Defects in Paired Eyes with Open-Angle Glaucoma. *PLoS One*. 2016;29(8):1-10.
  72. Yokoyama Y, Naoko A, Naoki C, Kazuko O, Masahiko N. Significant Correlations between Optic Nerve Head Microcirculation and Visual Field Defects and Nerve Fiber Layer Loss in Glaucoma Patient with Myopic Glaucomatous Disk. *Clinical Ophthalmology*. 2011;15(10):1721-7.
  73. Naito T, Keiji Y, Shiro M, Mami N, Tairo K. Relationship between Visual Field Progression and Baseline Refraction in Primary Open-Angle Glaucoma. *Clinical Ophthalmology*. 2016;10(12):197-1403.
  74. Ibanez MS, Jose MM, Gama R, Pilar CC, Maria NM, Laura MF. Utility of Bruch Membrane Opening-Based Optic Nerve Head Parameters in Myopic Subjects. *European Journal of Ophthalmology*. 2018;28(1):42-6.
  75. Lee YA, Yung FS, Luke LKL, Yehn YH, Tsing HW. Association Between High Myopia and Progression of Visual Field Loss in Primary Open-Angle Glaucoma. *J Formos Med Assoc*. 2008;107(12):952-7.
  76. Moroi H, Ayako A, Kyoko I, Goji T. Factors Related to a Right-Left Difference in The Eyes with Untreated Normal Tension Glaucoma. *Journal of Ophthalmology*. 2018;15(2):1-7.
  77. Hung CH, Shwu HL, Szu YL, Shung LL, Yi CC. The Relationship between Optic Nerve Head Deformation and Visual Field Defects in Myopic Eyes with Primary Open-Angle Glaucoma. *PLoS One*. 2018;13(12):1-14.
  78. Perera SA, Tien YW, Wan TT, Paul JF, Seang MS. Refractive Error, Axial Dimensions, and Primary Open-Angle Glaucoma. *Arch Ophthalmol*. 2010;128(7):900-5.
  79. Lucy KA, Gadi W. Structural and Functional Evaluations for The Early Detection of Glaucoma. *Expert Rev Ophthalmol*. 2016;11(5):367-76.
  80. Angraini R, Hendriati, Fitratul I. Hubungan Miopia dengan Ketebalan *Retinal Ganglion Cells* pada Mahasiswa Kedokteran. *Universitas Andalas*. 2017.
  81. Mahdavi KN, Nowroozizadeh S, Nassiri N, Cirineo C, Knipping S. Macular Ganglion Cell/Inner Plexiform Layer Measurements with Spectral Domain Optical Tomography for Detection of Early Glaucoma And Comparison to Retinal Nerve Fiber Layer Measurements. *Am J Ophthalmol*. 2013;12(6):1297-1307.
  82. Mwanza JC, Budenz D, Godfrey D, Neelakantan A, Sayyad F. Diagnostic Performance of Optical Coherence Tomography Ganglion Cell-Inner Plexiform Layer Thickness Measurements in Early Glaucoma. *Ophthalmology*. 2014;4(4):849-54.