

## DAFTAR PUSTAKA

1. Fung F, Sherrard DJ, Gillen DL, Wong C, Kestenbaum B, Ball A, et al. Increased risk for cardiovascular mortality among malnourished end stage renal disease patients. *Am J Kidney Dis.* 2002;40:307–14.
2. The National Kidney Foundation produces clinical practice guidelines through the NKF Kidney Disease Outcomes Quality Initiative (NKF-KDOQI). 2012.
3. Singh AK, Faragh YM, Mittal BV, Subramanian KK, Reddy SR, Acharya VN, et al. Epidemiology and risk factors of chronic kidney disease in India; results from the SEEK (Screening and Early Evaluation of Kidney Disease) study. *BMC Nephrology.* 2013;14(114):1–10.
4. Hill NR, Fatoba ST, Oke JL, Hirst JA, Callaghan CA, Lessersens DS, et al. Global prevalence of chronic kidney disease- a systematic review and meta-analysis. *Plos One.* 2016; 11(7):1–9
5. TIM Indonesia Renal Registry. Indonesian renal registry 10th annual report of indonesian renal registry. *Infodatin Kemenkes RI.* 2017;1–40.
6. Lucky VA, Toneeli N, Stanifersas. The Global burden of kidney disease and the sustainable development goals. *Bulletin of the World Health Organization* 2018;96 (6):414–22D.
7. Vaziri ND, Pahl MV, Crum A, Norris K. Effect of uremia on structure and function of immune system. *J Ren Nutr.* 2012;22(1):149–56.
8. Kato S, Chmielewski M, Honda H, Pecoits-Filho R, Matsuo S, Yuzawa Y et al. Aspects of immune dysfunction in end-stage renal disease. *Clin J Am Soc Nephrol.* 2008;3:1526–33.
9. Dalrymple LS, Go AS. Epidemiology of acute infections among patients with chronic kidney disease. *Clin J Am Soc Nephrol.* 2008;3:1487–93.
10. Beddhu S, Bruns FJ, Saul M. A simple comorbidity scale Predicts clinical outcomes and costs in dialysis patients. *American Journal Medicine* 2000;108(8):609–13.
11. Cobo G, Lindholm B, Stenvinkel P. Chronic inflammation in end-stage renal disease and dialysis. *Nefrology Dialysis Trasplant Journal.* 2018; 33(3): 35–40.
12. Ohno Y, Nakashima J, Ohori M, Hatano T, Tachibana M. Pretreatment neutrophil-to-lymphocyte ratio as an independent predictor of recurrence in patients with nonmetastatic renal cell carcinoma. *J Urol.* 2010;184:873–78.
13. Sarnak MJ, Jaber BL. Mortality caused by sepsis in patients with end-stage renal disease compared with the general population. *Kidney Int.* 2000; 58:1758–64.

14. Collins AJ, Foley RN, Herzog C, Gilbertson D, Chen S, Ishani A et al. Excerpts from the US renal data system 2009 annual data report. *Am J Kidney Disease*. 2010; 55:S1–A7.
15. Rojas L, Muñoz P, Kestler M, Bunsow E, Saez E, Sanchez C et al. Bloodstream infections in patients with kidney disease: risk factors for poor outcome and mortality. *J Hosp Infect* 2013; 85:196–205.
16. An X, Mao HP, Wei X, Chen JH, Guo N, Yang X, et al. Elevated neutrophil to lymphocyte ratio predicts overall and cardiovascular mortality in maintenance peritoneal dialysis patients. *Int Urol Nephrol*. 2012;44:1521–28.
17. Gulay O, Inal S, Erdem R, Ozge T, Esey Oyar, Guza Galip et al. Neutrophil to lymphocyte ratio of inflammation patient with Chronic Kidney Disease. *Informa Healthcare USA*. 2012;8 :1–8.
18. Erdem E. Neutrophil lymphocyte ratio in acute renal failure. *Indian J Nephrol*. 2015; 25(2): 126–27.
19. Wang Zebin, Wei Shi, Liang Jianbo, Liang Wejian, Liang Xinling. Association of interleukin 17 / angiotensin II with refractory hypertension risk in hemodialysis patients. *Afr Health Sci*. 2016 ; 16(3): 766–71.
20. Kitching AR, Ashley S, Stephen R, Shaun A, Summers X. Targeting IL-17 and IL-23 in immune mediated renal disease. *Curr Med Chem*. 2015;22(38):4341–65.
21. Mccharty M, Cooke K, Aronof DM, McMillan TR, Olkowitz K. IL-17 contributes to neutrophil recruitment but not to control of viral replication during acute mouse adenovirus type 1 respiratory Infection. *Virology*. 2014;5:259–67
22. Umani Vidia, Lidya Aidia, Nainggolan Gionova, Setiati Seti. Pengembangan model prediksi mortalitas 3 bulan pertama pada pasien penyakit ginjal kronik yang menjalani hemodialisa. *Jurnal Ilmu Penyakit Dalam FKUI*. 2000;2(3):170–82
23. Betjes MG. Immune cell dysfunction and inflammation in end-stage renal disease. *Nephrol*. 2013;9:255–65.
24. Puspa, Yunika Dewi. Performa formula Cockcroft–gault, MDRD dan CKD-EPI. *Patologi Klinik*. FK UGM. 2014.
25. Nafira Gani, Ali RH, Paat B. USG ginjal pada pasien PGK pada SMF Radiologi RSUP Kandou. *Jurnal e-clinic*. 2017;5(2):133–36.
26. Sudoyo AW, Setiyohadi B, Alwi I, K MS, Setiati S. *Buku ajar ilmu Penyakit Dalam*. I ed. Jakarta: Pusat Penerbitan Ilmu Penyakit Dalam FKUI; 2006. P. 570–3.

27. Brunner FP, Suddarth NH., 2001. End-Stage Renal failure due to analgesic nephropathy, its changing pattern and cardiovascular mortality. *Nephrol Dial Transplant*.2011; 9: 1371-76
28. Suharyanto, Majid. *Buku Panduan Hemodialisa* ; EGC.2009.Jakarta
29. Lindley E, Aspinal L, Gardiner C, Garthwaite E. Management of fluid status in hemodialysis patients ; *The Roles Of Technologi And Dietary Advice*.Leeds Teaching Hospital.2011:78-79
30. Cahyaning ND. *Hemodialisis panduan praktis perawatan gagal ginjal*. Mitra Cendekia. 2009.
31. Hoffbrand A. Victor. *Hematology at glance*. Edisi 6. Jakarta .EGC;2013;25-31.
32. Cendoroglo M, Jaber BL, Balakrishnan VS, Perianayagam M, King AJ, Pereira BJ. Neutrophil apoptosis and dysfunction in uremia. *J Am Soc Nephrol*. 1999;10:93-100.
33. Majewska E, Baj Z, Sulowska Z, Rysz J, Luciak M. Effects of uremia and haemodialysis on neutrophil apoptosis and expression of apoptosis-related proteins. *Nephrology Dialysis. Transplant*. 2003;18:2582-8.
34. Holub M, Beran O, Kasprikova N, Chalupa P. Neutrophil to lymphocyte count ratio as a biomarker of bacterial infections. *Cent Eur J Med* 2012;7:258-61.
35. Zahorec R. Ratio of neutrophil to lymphocyte count-rapid and simple parameter of systemic inflammation and stress in critically ill. *Bratis Lek Listy*. 2001;102:5-14.
36. Forget Patrice, Khalifa Celine, Jena Phelippe, Defour J, Latiene D, Pel MV, et al. What is the normal value of the neutrofil to lymphocyte. 2017. *BMC Research Notes*.2017; 10(12):1-4
37. Malaspina A, Moir S, Chaitt DG, Rehm CA, Kottlil S, Falloon J, et al. Idiopathic CD4+ T lymphocytopenia is associated with increases in immature/transitional B cells and serum levels of IL-7. *Blood*.2007: 109 (5): 2086-8.
38. Palmer MT, Weaver CT. Autoimmunity: increasing suspects in the CD4+ T. *Nat Immunol*. 2010; 11: 36-40.
39. Paust HJ, Turner JE, Steinmetz OM, Peters A, Heyman F, Holscher C, et al. The Th17 axis contributes to renal injury in experimental glomerulonephritis. *J Am Soc Nephrol*. 2009; 20: 969-79.
40. Summers SA, Steinmetz OM, Li M, Kaufman J, Serupie T, Edghon KL, et al. Th1 and Th17 cells induce proliferative glomerulonephritis. *J Am Soc Nephrol*. 2009; 20: 2518-24.

41. Gan PY, Ooi JD, Sullivan KM, Steinmetz OM, Tan DS, Summers SA, et al. Th 17 cells promote autoimmune anti-myeloperoxidase glomerulonephritis. *J Am Soc Nephrol*.2010;21(6):925–31
42. Loong CC, Hsieh HG, Lui WY, Chen A, Lyn CY. Evidence for the early involvement of interleukin 17 in human and experimental renal allograft rejection. *J Pathol*. 2002; 197: 322–32.
43. Crispin JC, Oukka M, Bayliss G. Expanded double negative T cells in patients with systemic lupus erythematosus produce IL-17 and infiltrate the kidneys. *J Immunol*. 2008; 181: 8761–66.
44. Wang Y, Ito S, Chino Y, Goto D, Matsumoto F, Hirata H, et al. Laser microdissection-based analysis of cytokine balance in the kidneys of patients with lupus nephritis. *Clin Exp Immunol*. 2010; 159: 1–10.
45. Yazawa Takasi, Shibata Mashahiko, Kenji Gonda, Machida T, Suzuki S,. Increased IL-17 production correlates with immunosuppression involving myeloid-derived suppressor cells and nutritional impairment in patients with various gastrointestinal cancers. *Clinical oncology journal*. 2013; 1(4): 675–79.
46. Zhang, Hai F, Yan L, Wang, Hong Y, Qian Li et al. Neutrophil to lymphocyte Ratio improves the accuracy and sensitivity of pneumonia severity index in predicting 30 day mortality of CAP patient. *Clin Lab Journal*.2019;65(10):1–9
47. Prasetya Dedi, Gunadi G, Rochadi R. Accuracy of neutrophil to lymphocyte ratio for diagnosis acute appendicitis; a diagnostic study. *Annals of Medicine and surgery Journal*. 2019;48:35–38
48. Collins A, Foley R Chavers B, Gilberton B, Herzog C, Shani A, et al. U.S. Renal Data System, *USRDS 2010 Annual Data Report*.2010;1:1–16
49. Webster AC, Nagler EV, Morton RL, Masson P. Chronic kidney disease. *The Lancet*.2017;389:1238–52.
50. Harford R, Clark MJ, Norris KC, Yan G. Relationship between age and pre-end stage renal disease care in elderly patients treated with maintenance hemodialysis. *Nephrol Nurs J*. 2016;43(2):101–8
51. Yositomi Ryota, Nakayama Nasaru, Kitazono Takanari. High neutrophil/lymphocyte ratio is associated with poor renal outcomes in Japanese patients with chronic kidney disease. *Renal Failure Journal*.2019;41(1):238–43
52. Marino Claudia, Feraro Pietro, Bargagli Mateo, Casini Silvia, Agabiti Nera, Gambaro Giovanni, et al. Prevalence of chronic kidney disease in the Lazio region, Italy: a classification algorithm based on health information systems. *BMC Nephrology*.2020;21(23):1–10

53. Elvira Dwitya. Korelasi Kadar Leukosit dan Rasio Netrofil-Limfosit dengan Kadar Interleukin- 17 pada Subjek Geriatri Penderita Pneumonia Komuniti. Bagian Ilmu Penyakit Dalam RSUP dr. M. Djamil Padang.
54. Sato Yu, Gonda Kenji, Yazawa T, Shibata Mashahito, Tanasaka Yuki, Sato Hiroshi, et al. Increased neutrophil-to-lymphocyte ratio is a novel marker for nutrition, inflammation and chemotherapy outcome in patients with locally advanced and metastatic esophageal squamous cell carcinoma. *Biomed Reports*.2017;1(1):79–84
55. Kaya T, Acikgoz SB, Yildirim M, Nalban A, Atlas AE, Cinemre H. Association between neutrophil-to-lymphocyte ratio and nutritional status in geriatric patients. *Journal of Clinical Analysis*.2019;33(1):1–10
56. Azikin Asriyani, Mangarengi Fitriani, Bahriun Uleng. Relationship between neutrophil to lymphocyte ratio and decreased glomerular filtration rate in diabetic. *Indonesian Journal of Clinical Pathology and Medical Laboratory*.2018;24(2):131–5
57. Turkmen K, Erdur FM, Ozcicek FM, Ozcicek A, Akbas EM, Ozbizer A et al. Platelet-to-lymphocyte ratio better predicts inflammation than neutrophil-to-lymphocyte ratio in end-stage renal disease patients. *Hemodialysis Internasional Journal*.2013;17(3):391–6
58. Catabay Christina, Obi Yoshitugu, Zadeh Kamyar. Lymphocyte cell ratios and mortality among incident hemodialysis patients. *American Journal of Nephrology*.2017;46(5):408–16
59. Chung BH, Kim KW, Sun IO, Choi SR, Park HS, Jeon NJ, et al. Increased interleukin-17 producing effector memory T cells in the end-stage renal disease patients. *Immunology Letters*.2012;141(2):181–9
60. Onishi RM, Gaffen SL. Interleukin-17 and its target genes: mechanisms of interleukin-17 function in disease. *The Journal of Cells, Molecules, System and Technologies*.2010;129(3):311–21
61. Kramer Stephanie, Loof Tanja, Harm Peters. Interleukin-17 and its target genes: mechanisms of interleukin-17 function in disease. *Plos One*.2016;15(5):1–19
62. Ghali JR, Holdsworth SR, Kitching AR. Targeting IL-17 and IL-23 in immune mediated renal disease. *Current Medical Chemistry*.2015;22(38):4341–65
63. Nelly , Widianingsih Yuyun, Mangarengi Fitriani. Nilai rasio netrofil limfosit dan rasio trombosit limfosit pada pasien chronic kidney disease sebelum dan setelah hemodialisa. *Inti sari sains Medis*.2019;10(2):230–4
64. Malik Ali Osman, Maowea Muhammad Mukhtar, Khalid husein Bhakeit, Zaki Hamdan. Plasma level Interlekin 17, interleukin 23 dan transforming growth factor B; A case control study. *Ijstr journal*.2015;60(6):634–9

65. Sharaawy Abdelbasiit, Halim Aber Baki, Nahla Mohammed, Rania Abdel, Alaa Noha, Ahmed Reem. Neutrophils to Lymphocytes ratio is an easy non expensive marker of inflammation in hemodialysis. *Journal of Clinical & Experimental Nephrology*.2018;2(4):1–6
66. Yu X. The evolving patterns of uremia: unmet clinical needs in dialysis. *Contrib Nephrol* .2017; 19(1): 1–7
67. Zenobia Camille, Haesengalis George. Basic biology and role of interleukin-17 in immunity and inflammation. *Periodontolgy Journal*. 2000;69(2):72–6
68. Anding Kristen, Gross Ptere, M jan, Allegair Dirk, Jacobs Enno. The influence of uraemia and haemodialysis on neutrophil phagocytosis and antimicrobial killing. *Nephrology Dyalysis Transpalantation Journal*.2003;18(8): 2067–773
69. Shimamura Shunshuke, Kimura Ken, Yasuda Jun. Evaluation of neutrophil function during hemodialysis treatment in healthy dogs under anesthesia with sevoflurane. *Journal Vet Medical Science*.2014;76(11): 1539–43
70. Peres Allesandra, Perotto Lorenzo Daniella, Dorneles Gilson, Isabel Maria, Monteiro Mariane. Effects of intradialytic exercise on systemic cytokine in patients with chronic kidney disease. *Renal Failure Journal*. 2015; 37(9) :1430–4
71. Mandal K Anil, Sun Yuuan, Servilas S Karen, Antonius H. *Text Book of Nephrology* 3rd edition.2014. Jaypee Brothers Medical Publishers:397
72. Liang Kelly, Jane H zhang, Valesky Paul. Urea reduction ratio may be a simpler approach for measurement of adequacy of intermittent hemodialysis in acute kidney injury. *BMC Nephrology*.2019;20(82):1272–7
73. Valeri Maria, Raffatellu Manuela. Cytokines IL-17 and IL-22 in the host response to infection. *Patoghen and Disease Journal*.2016:74:111
74. Hecking Manfred, Biebier Brian, Eather Jean, kautzky Alexandra, Plasman Gere, Sausman Marcus, et al. Sex-Specific Differences in Hemodialysis Prevalence and Practices and the Male-to-Female Mortality Rate: The Dialysis Outcomes and Practice Patterns Study (DOPPS). *Plos Medicine Journal*. 2014:11:10
75. Aisara Sitifa, Azmi Syaiful, Yanni Mefri. Gambaran klinis penderita penyakit ginjal kronik yang menjalani hemodialisis di RSUP dr. M. Djamil.2018. *Jurnal kesehatan Andalas*:7:1
76. Zyga S, Saraves P. Hemodialysis Adequacy. *Health Medical Journal*. 2009: 209–213

