


DAFTAR PUSTAKA

1. Sukandar EY, Andrajati R, Sigit JI, Adnyana IK, Setiadi AAP, Kusnandar. Iso Farmakoterapi. Jakarta: PT ISFI Penerbitan; 2008.
2. Kemenkes RI. Farmakope Indonesia edisi V. Jakarta: Direktorat Jendral Bina Kefarmasian dan Alat Kesehatan Republik Indonesia; 2014.
3. Moffat AC, Osselton MD, Widdop B. Clarke's Analysis of Drugs and Poisons 4th Ed. London: Pharmaceutical Press; 2011.
4. American Society of Health-System Pharmacists. AHFS Drug Information® Essentials. Bethesda: American Society of Health-System Pharmacists®; 2011.
5. Nikhade A, Mulgund S V. UV spectrophotometric estimation of diltiazem hydrochloride in bulk and tablet dosage form using area under curve method. World J Pharm Pharm Sci. 2014;3(8):1217–24.
6. Hassan A, El-asmy AF, El-Raheem YBA. Sensitive extractive simpler spectrophotometric methods for the microdetermination of diltiazem HCL and pipazethate HCL in pure and tablet dosage forms. Asian J Res Chem. 2011;4(9):1489–502.
7. Patil R, Patil V, Bhosale S. Development and validation of zero order derivative by uv- spectrophotometric method for determination of diltiazem hcl in bulk and its formulation. Int J Univers Pharm bio Sci [Internet]. 2014;3(4):1–12. Available from: <http://www.ijupbs.com/Uploads/1.RPA1415104015.pdf>
8. Sabino LZL, Marino DC, Moya HD. Determination of diltiazem based on

the reduction of Cu (II)– BCA complexes in micellar medium. Can J Chem [Internet]. 2010;88:533–9. Available from: <https://doi.org/10.1139/V10-036>

9. Ayad MM, Abdellatef HE, Hosny MM, Sharaf YA. Application of 4-chloro-7-nitrobenzofurazan for the analysis of propafenone and diltiazem hydrochlorides using kinetic spectrophotometric and spectrofluorimetric methods. Eur J Chem [Internet]. 2013;4(1):35-43. Available from: <https://doi.org/10.5155/eurjchem.4.1.35-43.713>
10. Hassan AM, F.El-Asmy A, El-Raheem YBA. Novel methods for the visible spectrophotometric determination of diltiazem HCl and levamisole HCl in pure, tablet dosage forms and biological fluids. Anal Chem an indian J. 2011;10(11):740–53.
11. Omar MA, Abdelmageed OH, Abdelgaber AA, Saleh SF. Spectrophotometric determination of some calcium channel blockers using Sulfochlorophenol-S , Pyrocatechol violet. Nat Sci [Internet]. 2013;5(4):514–25. Available from: <http://dx.doi.org/10.4236/ns.2013.54065>
12. Shrivastava R. Spectrophotometric determination of diltiazem in pharmaceutical and in-vivo samples with Pd(II). Int J Pharm Sci Res [Internet]. 2013;4(12):4676–81. Available from: [http://dx.doi.org/10.13040/IJPSR.0975-8232.4\(12\).4676-81](http://dx.doi.org/10.13040/IJPSR.0975-8232.4(12).4676-81)
13. Sultana N, Arayne MS, Shafi N, Siddiqui FA, Hussain A. Development of a RP-HPLC method for the simultaneous analysis of diltiazem and statin: Application in pharmaceuticals and human serum. Anal Methods [Internet]. 2010;2:1571–6. Available from: <https://doi.org/10.1039/C0AY00337A>

14. Sultana N, Arayne MS, Shafi N, Siddiqui FA. Simultaneous RP-LC analysis of diltiazem and non-steroidal anti-inflammatory drugs in pharmaceutical formulations and human serum. *Chromatographia*. 2010;71(1/2):71–7.
15. Sultana N, Arayne MS, Shafi N, Siddiqui FA, Hussain A. Development and validation of new assay method for the simultaneous analysis of diltiazem, metformin, pioglitazone and rosiglitazone by RP-HPLC and its applications in pharmaceuticals and human serum. *J Chromatogr Sci* [Internet]. 2011;49:774–9. Available from: <https://doi.org/10.1093/chrscl/49.10.774>
16. Chatpalliwar VA, Porwal PK, Upmanyu N. Validated gradient stability indicating HPLC method for determining Diltiazem Hydrochloride and related substances in bulk drug and novel tablet formulation. *J Pharm Anal* [Internet]. 2012;2(3):226–37. Available from: <http://dx.doi.org/10.1016/j.jpha.2012.01.003>
17. Abu-nameh ESM. A validated stability indicating HPLC method for determination of diltiazem hydrochloride in tablet dosage form. *Aust J Basic Appl Sci*. 2013;7(6):730–6.
18. Sadeghi F, Navidpour L, Bayat S, Afshar M. Validation and uncertainty estimation of an ecofriendly and stability-indicating hplc method for determination of diltiazem in pharmaceutical preparations. *J of Analytical Methods Chem* [Internet]. 2013;1–10. Available from: <http://dx.doi.org/10.1155/2013/353814>
19. Li N, Li C, Lu N, Dong Y. A novel micellar per aqueous liquid chromatographic method for simultaneous determination of diltiazem hydrochloride , metoprolol tartrate and isosorbide mononitrate in human serum. *J Chromatogr B* 967 [Internet]. 2014;90–7. Available from:



<http://dx.doi.org/10.1016/j.jchromb.2014.07.019>

20. Arafat M. Simple HPLC validated method for the determination of diltiazem hydrochloride in human plasma. *Int J Pharm Pharm Sci.* 2014;6(9):213–6.
21. Taylor P, Shafi N, Siddiqui FA, Sultana N, Arayne MS. Concurrent determination of diltiazem, lisinopril, captopril and enalapril in dosage formulations and in human serum by liquid chromatographic technique. *J Liq Chromatogr Relat Technol* [Internet]. 2015; Available from: <http://dx.doi.org/10.1080/10826076.2015.1050503>
22. Kumar BMS, Rajkamal B, Chandramowli B. A validated RP-HPLC method for the determination of diltiazem in raw Material and pharmaceutical dosage form. *Int J Pharm Sci Drug Res* [Internet]. 2018;10(6):487–91. Available from: <https://doi.org/10.25004/IJPSDR.2018.100609>
23. Naz S. Simultaneous determination of prazosin and calcium channel blockers in raw materials, pharmaceutical formulations and human serum by RP-HPLC. *Int J Pharma Res Dev.* 2010;2(9):6–12.
24. Soltani S, Jouyban A. Optimization and validation of an isocratic HPLC-UV method for the simultaneous determination of five drugs used in combined cardiovascular therapy in human plasma. *Asian J Chem.* 2011;23(4):1728–34.
25. Kulkarni AS, Jadhav SD, Khetmar SS, Bhatia MS. Development of chromatographic technique for simultaneous estimation of lovastatin and diltiazem hydrochloride. *Mahidol Univ J Pharm Sci.* 2012;39(3–4):17–23.

26. Eggadi V, Ponna SK, Kankanala SR, Sheshagiri SBB, Gaddam SR. Determination of simvastatin and diltiazem in rat's plasma by HPLC and pharmacokinetic studies. *Int J Pharm Sci Rev Res*. 2013;21(1):53–6.
27. Sultana N, Arayne MS, Ali SN. An Ultra-sensitive and selective LC-UV method for the simultaneous determination of pravastatin , diltiazem , naproxen sodium and meloxicam in API , pharmaceutical formulations and human serum. *Am J Appl Chem* [Internet]. 2013;1(1):1–8. Available from: <https://doi.org/10.11648/j.ajac.20130101.11>
28. Komesli OT, Bakirdere S, Bayoren C, Gokcay CF. Simultaneous determination of selected endocrine disrupter compounds in wastewater samples in ultra trace levels using HPLC-ES-MS/MS. *Env Monit Assess* [Internet]. 2012;184:5215–24. Available from: <https://doi.org/10.1007/s10661-011-2334-x>
29. Muz M, Snmez MS, Komesli OT, Bakirdere S, Gokcay CF. Determination of selected natural hormones and endocrine disrupting compounds in domestic wastewater treatment plants by liquid chromatography electrospray ionization tandem mass spectrometry after solid phase extraction. *Analist*. 2012;137:884–9.
30. Sultana N, M SA, Shah SN. Development and validation for the simultaneous quantification of prazosin, amlodipine, diltiazem and verapamil in API, dosage formulation and human serum by RP-HPLC : application to in vitro interaction studies. *Med Chem (Los Angeles)* [Internet]. 2014;4(12):770–7. Available from: <http://dx.doi.org/10.4172/2161-0444.1000228>
31. Patil BR, O.G B, N PB, Y GA, Mulaje SS. Analytical method development and validation for the estimation of diltiazem hydrochloride in bulk and

pharmaceutical dosage form by RP-HPLC. *Int J Drug Regul Aff.* 2014;2(2):78–84.

32. Pontarolo R, Floriani G, Gonc AG. Development and validation of an HPLC-DAD method for simultaneous determination of cocaine , benzoic acid , benzoylecgonine and the main adulterants found in products based on cocaine. *Forensic Sci Int* [Internet]. 2014;235:32–9. Available from: <http://dx.doi.org/10.1016/j.forsciint.2013.11.013>
33. Jouyban A, Sorouraddin MH, Farajzadeh MA, Somi MH, Fazeli-Bakhtiyari R. Determination of five antiarrhythmic drugs in Human plasma by dispersive liquid–liquid microextraction and High-performance liquid chromatography. *Talanta* [Internet]. 2014; Available from: <http://dx.doi.org/10.1016/j.talanta.2014.12.008>
34. Sena LCS, Matos HR, Dórea HS, Pimentel MF, Santana DCAS de, Santana FJM de. Dispersive liquid-liquid microextraction based on solidification of floating organic drop and high-performance liquid chromatography to the analysis of cocaine ' s major adulterants in human urine. *Toxicology* [Internet]. 2015; Available from: <http://dx.doi.org/10.1016/j.tox.2016.04.008>
35. Sonmez MS, Muz M, Komesli OT, Bakirdere S, Gokcay CF. Determination of selected endocrine disrupter compounds at trace levels in sewage sludge samples. *Clean – Soil, Air, Water.* 2012;00(0):1–6.
36. Catarino RIL, Leal MFC, Pimenta AM, Souto MRS, Lopes JRT. Cathodic voltammetric detection of diltiazem at a bismuth film electrode: Application to human urine and pharmaceuticals. *J Braz Chem Soc* [Internet]. 2014;25(5):961–8. Available from: <http://dx.doi.org/10.5935/0103-5053.20140067 J>.

37. Hasanzadeh M, Pournaghi-Azar MH, Shadjou N. Determination of Diltiazem in the presence of Timolol in human serum samples using a nanoFe₃O₄@GO modified glassy carbon electrode. *R Soc Chem Adv* [Internet]. 2014;4:51734–44. Available from: <https://doi.org/10.1039/C4RA05989A>
38. Gevaerd A, Caetano FR, Oliveira PR, Zarbin AJG, Bergamini MF, Marcolino-Junior LH. Thiol-capped gold nanoparticles: Influence of capping amount on electrochemical behavior and potential application as voltammetric sensor for diltiazem. *Sensors Actuators, B Chem* [Internet]. 2015;220:673–8. Available from: <http://dx.doi.org/10.1016/j.snb.2015.06.010>
39. Attaran AM, Abdol-Manafi S, Javanbakht M, Enhessari M. Voltammetric sensor based on Co₃O₄/SnO₂ nanopowders for determination of diltiazem in tablets and biological fluids. *J Nanostructure Chem* [Internet]. 2016;6:121–8. Available from: <https://doi.org/10.1007/s40097-015-0186-6>
40. Salamanca-Neto CAR, Felsner ML, Galli A, Sartori ER. In-house validation of a totally aqueous voltammetric method for determination of diltiazem hydrochloride. *J Electroanal Chem* [Internet]. 2019;837:159–66. Available from: <https://doi.org/10.1016/j.jelechem.2019.02.026>
41. Elfiky M, Salahuddin N, Matsuda A. Green fabrication of 3D hierarchical blossom-like hybrid of peeled montmorillonite-ZnO for in-vitro electrochemical sensing of diltiazem hydrochloride drug. *Mater Sci Eng C* [Internet]. 2020;111:110773. Available from: <https://doi.org/10.1016/j.msec.2020.110773>
42. Deng B, Lu H, Li L, Shi A, Kang Y, Xu Q. Determination of the number of binding sites and binding constant between diltiazem hydrochloride and

human serum albumin by ultrasonic microdialysis coupled with online capillary electrophoresis electrochemiluminescence. *J Chromatogr A* [Internet]. 2010;1217:4753–6. Available from: <http://dx.doi.org/10.1016/j.chroma.2010.05.021>

43. Sanchez MA, Rocha FRP. Liquid-liquid microextraction without phase separation in a multicommuted flow system for diltiazem determination in pharmaceuticals. *Anal Chim Acta* [Internet]. 2011;694:95–9. Available from: <http://dx.doi.org/10.1016/j.aca.2011.03.051>

