

## DAFTAR PUSTAKA

1. Swanson CA, Burns JC, Peterson BM. Low plasma d-dimer concentration predicts the absence of traumatic brain injury in children. *J Trauma - Inj Infect Crit Care.* 2010;68(5):1072–7.
2. Langlois JA, Rutland-Brown W, Wald MM. The epidemiology and impact of traumatic brain injury: A brief overview. *J Head Trauma Rehabil.* 2006;21(5):375–8.
3. Corrigan JD, Selassie AW, Orman JA. The epidemiology of traumatic brain injury. *J Head Trauma Rehabil.* 2010;25(2):72–80.
4. Kementerian Kesehatan Republik Indonesia. No Title [Internet]. Hasil utama RISKESDAS 2018. 2018. Available from: <http://www.kemkes.go.id/resources/download/info-terkini/hasil-riskesdas-2018>
5. Bales JW, Bonow RH, Ellenbogen RG. Closed Head Injury [Internet]. Fourth Edi. Principles of Neurological Surgery. Elsevier Inc.; 2018. 366-389.e4 p. Available from: <https://doi.org/10.1016/B978-0-323-43140-8.00025-1>
6. Saanin S. Cedera Otak Traumatik [Internet]. 2010. Available from: <http://www.angelfire.com/nc/neurosurgery/Klasifikasi.html>
7. Young RJ, Destian S. Imaging of traumatic intracranial hemorrhage. *Neuroimaging Clin N Am.* 2002;12(2):189–204.
8. Yuan F, Ding J, Chen H, Guo Y, Wang G, Gao WW, et al. Predicting progressive hemorrhagic injury after traumatic brain injury: Derivation and validation of a risk score based on admission characteristics. *J Neurotrauma.* 2012;29(12):2137–42.
9. Liu J, Tian HL. Relationship between trauma-induced coagulopathy and progressive hemorrhagic injury in patients with traumatic brain injury. *Chinese J Traumatol - English Ed [Internet].* 2016;19(3):172–5. Available from: <http://dx.doi.org/10.1016/j.cjtee.2016.01.011>
10. Arifin M., Risdianto A. Cedera Kepala. Jakarta, Indonesia: Sagung Seto; 2013. 53–58 p.
11. Delgado P, Álvarez-Sabín J, Abilleira S, Santamarina E, Purroy F, Arenillas JF, et

- al. Plasma D-dimer predicts poor outcome after acute intracerebral hemorrhage. *Neurology*. 2006;67(1):94–8.
12. Subedi PP, Shilpkar SK. Correlation of D-Dimer level with outcome in traumatic brain injury. *J Soc Surg Nepal*. 2016;17(1):21–30.
13. Xu DX, Du WT, Li X, Wu ZX, Yu GF. D-dimer/fibrinogen ratio for the prediction of progressive hemorrhagic injury after traumatic brain injury. *Clin Chim Acta* [Internet]. 2020;507(April):143–8. Available from: <https://doi.org/10.1016/j.cca.2020.04.022>
14. Amin Foaud HM, Labib JR, Gabr Metwally H, Abd El-Twab KM. Plasma D-dimer as a prognostic marker in ICU admitted Egyptian children with traumatic brain injury. *J Clin Diagnostic Res*. 2014;8(9):PC01–5.
15. Hayakawa M, Maekawa K, Kushimoto S, Kato H, Sasaki J, Ogura H, et al. Hyperfibrinolysis in severe isolated traumatic brain injury may occur without tissue hypoperfusion: A retrospective observational multicentre study. *Crit Care*. 2017;21(1):1–9.
16. Sharma A, Sikka M, Gomber S, Sharma S. Plasma fibrinogen and D-dimer in children with sepsis: A single-center experience. *Iran J Pathol*. 2018;13(2):272–5.
17. Yeatts SD, Palesch YY, Temkin N. Biostatistical Issues in TBI Clinical Trials [Internet]. Second Edi. *Handbook of Neuroemergency Clinical Trials: Second Edition*. Elsevier Inc.; 2018. 167–185 p. Available from: <https://doi.org/10.1016/B978-0-12-804064-5.00009-6>
18. Callahan BP, Rabb CH. Traumatic Brain Injury [Internet]. *Abernathy's Surgical Secrets*. Elsevier Inc.; 2009. 99–104 p. Available from: <http://dx.doi.org/10.1016/B978-0-12-648082-5.50007-1>
19. Galgano M, Toshkezi G, Qiu X, Russell T, Chin L, Zhao LR. Traumatic brain injury: Current treatment strategies and future endeavors. *Cell Transplant*. 2017;26(7):1118–30.
20. Maegele M. Coagulopathy after traumatic brain injury: Incidence, pathogenesis, and treatment options. *Transfusion*. 2013;53(SUPPL. 1).
21. Saatman KE, Duhaime AC, Bullock R, Maas AIR, Valadka A, Manley GT, et al. Classification of traumatic brain injury for targeted therapies. *J Neurotrauma*.

- 2008;25(7):719–38.
22. Dash HH, Chavali S. Management of Traumatic Brain Injury Patient. Korean J Anesthesiol [Internet]. 2018;12–21. Available from: <http://ekja.org>
  23. Carney N, Totten AM, O'Reilly C, Ullman JS, Hawryluk GWJ, Bell MJ, et al. Guidelines for the Management of Severe Traumatic Brain Injury, Fourth Edition. Neurosurgery. 2017;80(1):6–15.
  24. American College of Surgeons. BEST PRACTICES IN THE MANAGEMENT OF TRAUMATIC Table of Contents. 2015;(January):1–29. Available from: [https://www.facs.org/~media/files/quality\\_programs/trauma/tqip/traumatic\\_brain\\_injury\\_guidelines.ashx](https://www.facs.org/~media/files/quality_programs/trauma/tqip/traumatic_brain_injury_guidelines.ashx)
  25. Capizzi A, Woo J, Verduzco-Gutierrez M. Traumatic Brain Injury: An Overview of Epidemiology, Pathophysiology, and Medical Management. Med Clin North Am [Internet]. 2020;104(2):213–38. Available from: <https://doi.org/10.1016/j.mcna.2019.11.001>
  26. Senn R, Elkind MSV, Montaner J, Christ-Crain M, Katan M. Potential role of blood biomarkers in the management of nontraumatic intracerebral hemorrhage. Cerebrovasc Dis. 2014;38(6):395–409.
  27. Olson JD. D-dimer: An Overview of Hemostasis and Fibrinolysis, Assays, and Clinical Applications [Internet]. 1st ed. Vol. 69, Advances in Clinical Chemistry. Elsevier Inc.; 2015. 1–46 p. Available from: <http://dx.doi.org/10.1016/bs.acc.2014.12.001>
  28. Johnson ED, Schell JC, Rodgers GM. The D-dimer assay. Am J Hematol. 2019;94(7):833–9.
  29. Quinn DA, Fogel RB, Smith CD, Laposata M, Thompson BT, Johnson SM, et al. D-Dimers in the Diagnosis of Pulmonary Embolism. (7).
  30. Sugimoto K, Suehiro E, Shinoyama M, Sadahiro H, Haji K, Fujiyama Y, et al. D-Dimer Elevation as a Blood Biomarker for Detection of Structural Disorder in Mild Traumatic Brain Injury. J Neurotrauma. 2017;34(23):3245–8.
  31. Becker DM, Philbrick JT, Bachhuber TL, Humphries JE. D-dimer testing and acute venous thromboembolism: A shortcut to accurate diagnosis? Arch Intern Med. 1996;156(9):939–46.

32. Oliveira R., Araujo S. Glasgow outcome scale at hospital discharge as a prognostic index in patients with severe traumatic brain injury. *Arq Neuropsiquiatr* [Internet]. 2012;70(8):604–8. Available from: <http://www.scielo.br/pdf/anp/v70n8/v70n8a09.pdf> %5Cn<http://ovidsp.ovid.com/ovidweb.cgi?T=JS&PAGE=reference&D=emed10&NEWS=N&AN=2012499953>
33. Chhabra G, Sharma S, Subramanian A, Agrawal D, Sinha S, Mukhopadhyay AK. Coagulopathy as prognostic marker in acute traumatic brain injury. *J Emergencies, Trauma Shock.* 2013;6(3):180–5.
34. Kuo JR, Lin KC, Lu CL, Lin HJ, Wang CC, Chang CH. Correlation of a high D-dimer level with poor outcome in traumatic intracranial hemorrhage. *Eur J Neurol.* 2007;14(10):1073–8.
35. Chen S, Zhao B, Wang W, Shi L, Reis C, Zhang J. Predictors of hematoma expansion predictors after intracerebral hemorrhage. *Oncotarget.* 2017;8(51):89348–63.