

DAFTAR PUSTAKA

1. Kusumastuti DA, Taufiq N, Mumpuni H. correlation between Level of Soluble Suppression of Tumorigenicity-2(sST2) with Global Longitudinal Strain(GLS) of Left Ventricle inPatients with Acute Myocardial Infraction. *Acta Cardiol Indones*. 2019;5(1):62–70.
2. Pramadiatz AT, Fadil M, Mulyani H. Artikel Penelitian Hubungan Faktor Risiko Terhadap Kejadian Sindroma Koroner Akut pada Pasien Dewasa Muda di RSUP Dr . M . Djamil Padang. *J Kesehat Andalas*. 2016;5(2):330–7.
3. Faiqotun B, Herdianto D, Iskandar A. Gambaran Lokasi Lesi Culprit, Elektrokardiogram, dan TIMI Flow Pada Pasien IMA-EST yang telah dilakukan IKPP. *VerdureHealth Sci J*. 2019;1(1):1–12.
4. Perhimpunan Dokter Spesialis Kardiovaskular Indonesia. Pedoman Tatalaksana Sindrom Koroner Akut. *J Kardiologi Indones*. 4th ed. 2018.
5. Departemen kardiologi & kedokteran vaskular. Buku Ajar Kardiovaskular Jilid 2. Yuniadi Y, Hermanto DY, Siswanto BB, editors. Jakarta; 2017.
6. Loh JP, Tan LL, Zheng H, Lau YH, Chan SP, Tan KB, et al. First Medical Contact-to-Device Time and Heart Failure Outcomes Among Patients Undergoing Primary Percutaneous Coronary Intervention. *Circ Cardiovasc Qual Outcomes*. 2018;11(8).
7. Nepper-Christensen L, Lønborg J, Høfsten DE, Sadjadieh G, Schoos MM, Pedersen F, et al. Clinical outcome following late reperfusion with percutaneous coronary intervention in patients with ST-segment elevation myocardial infarction. *Eur Hear journal Acute Cardiovasc care*. 2020.
8. Khalfallah M, Elsheikh A, Abdalaal M. Very Early Versus Early Percutaneous Coronary Intervention After Successful Fibrinolytic Therapy in Pharmacoinvasive Strategy. *Glob Heart*. 2018;13(4):261–5.
9. Morshed MG, Azam M, Arefin MM, Hussain KS, Jahan J, Ahmed N, et al. Impact of Time Delay on Short Term Outcome of Primary Percutaneous Coronary Intervention (PCI) in Bangladeshi Population. *Univ Hear J*. 2019;14(2):77–82.
10. Martalena D. Pengaruh Hiperglikemia Admisi terhadap Major Adverse Cardiac Events Selama Perawatan pada Pasien Sindrom Koroner Akut di ICCU RSCM, Jakarta. *eJournal Kedokt Indones*. 2013;1(2):106–12.
11. Tsai IT, Wang CP, Lu YC, Hung WC, Wu CC, Lu LF, et al. The burden of major adverse cardiac events in patients with coronary artery disease. *BMC Cardiovasc Disord*. 2017;17(1):1–13.

12. Wilar GI, Panda AL, Rampengan SH. Pengamatan 6 Bulan Terhadap Kejadian Kardiovaskular Mayor Pada Pasien dengan Infark Miokard Akut dengan Elevasi Segmen ST (STEMI) di RSUP Prof. Dr.R.D. Kandou Manado Periode Januari - Desember 2017. *J Med dan Rehabil.* 2019;1(3):1–8.
13. Gao M, Qin L, Zhang Z, Chen L, Zheng Y, Tong Q, et al. Treatment Windows and Clinical Outcomes in Late-Presenting Patients with ST-Segment Elevation Myocardial Infarction. *Am J Med Sci.* 2019;358(4):248–55.
14. Chandrasekhar J, Marley P, Allada C, McGill D, O'Connor S, Rahman M, et al. Symptom-to-Balloon Time is a Strong Predictor of Adverse Events Following Primary Percutaneous Coronary Intervention: Results From the Australian Capital Territory PCI Registry. *Heart Lung Circ.* 2017;26(1):41–8.
15. Saleh M, Ambrose JA. Understanding myocardial infarction. *F1000Research.* 2018;7(0):1–8.
16. Ibanez B, James S, Agewall S, Antunes MJ, Bucciarelli-Ducci C, Bueno H, et al. 2017 ESC Guidelines for the management of acute myocardial infarction in patients presenting with ST-segment elevation. *Eur Heart J.* 2018;39(2):119–77.
17. Bulluck H, Zheng H, Chan MY, Foin N, Foo DC, Lee CW, et al. Independent Predictors of Cardiac Mortality and Hospitalization for Heart Failure in a Multi-Ethnic Asian ST-segment Elevation Myocardial Infarction Population Treated by Primary Percutaneous Coronary Intervention. *Sci Rep.* 2019;9(1):1–14.
18. Dharma S, Andriantoro H, Purnawan I, Dakota I, Basalamah F, Hartono B, et al. Characteristics, treatment and in-hospital outcomes of patients with STEMI in a metropolitan area of a developing country: an initial report of the extended Jakarta Acute Coronary Syndrome registry. *BMJ Open.* 2016;6(8).
19. PRARA MR. Kejadian Kardiovaskular Mayor Selama Rawatan Pada Infark Miokard Akut dengan Elevasi Segmen ST (IMA-EST) di RSUP Dr. M. Djamil Padang. Universitas Andalas; 2017.
20. Lilly LS. Patofisiologi penyakit jantung. Edisi 6. Juzar DA, Ridjab DA, Sari IP, Putratama R, editors. Jakarta: MEDIK; 2019.
21. Bentzon JF, Falk E. Pathogenesis of Stable and Acute Coronary Syndromes. In: *Acute coronary syndromes.* 2nd editio. United states of America: Elsevier saunders; 2011. p. 42–52.

22. Aluganti Narasimhulu C, Fernandez-Ruiz I, Selvarajan K, Jiang X, Sengupta B, Riad A, et al. Atherosclerosis - Do we know enough already to prevent it? *Curr Opin Pharmacol*. 2016;27(Ldl):92–102.
23. Asada Y, Yamashita A, Sato Y, Hatakeyama K. Thrombus formation and propagation in the onset of cardiovascular events. *J Atheroscler Thromb*. 2018;25(8):653–64.
24. Alwi I. Infark Miokard Akut dengan Elevasi ST. In: Setiati S, Alwi I, Sudoyo A., K MS, Setiyohadi B, Syam AF, editors. *Buku ajar Ilmu Penyakit Dalam Jilid II*. Edisi VI. Jakarta: Interna Publishing; 2014. p. 1459–76.
25. Stouffer G, Runge MS, Patterson C, Rossi JS. *Netter's Cardiology*. 3rd ed. Elsevier; 2018.
26. Reed GW, Rossi JE, Cannon CP. Acute myocardial infarction. *Lancet*. 2017;389(10065):197–210.
27. Vogel B, Claessen BE, Arnold S V., Chan D, Cohen DJ, Giannitsis E, et al. ST-segment elevation myocardial infarction. *Nat Rev Dis Prim*. 2019;5(1):1–20.
28. Qodir A. Kepatuhan Mengontrol Tekanan Darah Sebagai Upaya Menurunkan Kejadian Infark Miokard Akut Recurrent. *J Ilm Kesehat Media Husada*. 2016;5(1):11–8.
29. Taniguchi T, Shiomi H, Morimoto T, Watanabe H, Ono K, Shizuta S, et al. Incidence and Prognostic Impact of Heart Failure Hospitalization During Follow-Up After Primary Percutaneous Coronary Intervention in ST-Segment Elevation Myocardial Infarction. *Am J Cardiol*. 2017;119(11):1729–39.
30. Sacco RL, Kasner SE, Broderick JP, Caplan LR, Connors JJ, Culebras A, et al. An updated definition of stroke for the 21st century: A statement for healthcare professionals from the American heart association/American stroke association. *Stroke*. 2013;44(7):2064–89.
31. Keeley EC, Hillis LD. Primary PCI for myocardial infarction with ST-segment elevation. *N Engl J Med*. 2007;356(1):47–54.
32. RÁCZ I, FÜLÖP L, KOLOSVÁRI R, SZABÓ GT, BÓDI A, PÉTER A, et al. Wall motion changes in myocardial infarction in relation to the time elapsed from symptoms until revascularization. *Anadolu Kardiyol Derg*. 2015;15(5):363–70.
33. Wulan Sari NMA. Waktu antara angina dan tindakan percutaneous

coronary intervention menyebabkan perubahan wall motion score index pada pasien infark miokard akut. *Medicina (B Aires)*. 2018;49(3):314–9.

34. Scholz KH, Maier SKG, Maier LS, Lengenfelder B, Jacobshagen C, Jung J, et al. Impact of treatment delay on mortality in ST-segment elevation myocardial infarction (STEMI) patients presenting with and without haemodynamic instability: Results from the German prospective, multicentre FITT-STEMI trial. *Eur Heart J*. 2018;39(13):1065–74.
35. Shafi MJ, Nasrin S. Outcome of Early vs. Delayed Invasive Intervention in Acute Coronary Syndrome Patients Attending a Selected Specialized Hospital: A Comparative Study. *Cardiovasc J*. 2019;11(2):129–38.
36. Gerber RT, Arri SS, Mohamed MO, Dhillon G, Bandali A, Harding I, et al. Age is not a bar to PCI: Insights from the long-term outcomes from off-site PCI in a real-world setting. *J Interv Cardiol*. 2017;30(4):347–55.
37. Bauer T, Zeymer U. Impact of age on outcomes of percutaneous coronary intervention in acute coronary syndromes patients. *Interv Cardiol*. 2010;2(3):319–25.
38. Heer T, Hochadel M, Schmidt K, Mehilli J, Zahn R, Kuck KH, et al. Sex Differences in Percutaneous Coronary Intervention-Insights From the Coronary Angiography and PCI Registry of the German Society of Cardiology. *J Am Heart Assoc*. 2017;6(3).
39. Endo H, Dohi T, Shitara J, Wada H, Doi S, Naito R, et al. Impact of body mass index on long-term outcomes in Japanese patients following percutaneous coronary intervention: The Juntendo PCI Registry. *J Cardiol*. 2018;72(3):208–14.
40. Orbach A, Halon DA, Jaffe R, Rubinshtein R, Karkabi B, Flugelman MY, et al. Impact of diabetes and early revascularization on the need for late and repeat procedures. *Cardiovasc Diabetol*. 2018;17(1):1–9.
41. Saluveer O, Redfors B, Angerås O, Dworeck C, Haraldsson I, Ljungman C, et al. Hypertension is associated with increased mortality in patients with ischaemic heart disease after revascularization with percutaneous coronary intervention—a report from SCAAR*. *Blood Press*. 2017;26(3):166–73.
42. Reindl M, Holzkecht M, Tiller C, Lechner I, Schiestl M, Simma F, et al. Impact of infarct location and size on clinical outcome after ST-elevation myocardial infarction treated by primary percutaneous coronary intervention. *Int J Cardiol*. 2020;301:14–20.
43. Wang R, Mei B, Liao X, Lu X, Yan L, Lin M, et al. Determination of risk factors affecting the in-hospital prognosis of patients with acute ST segment elevation myocardial infarction after percutaneous coronary

intervention. *BMC Cardiovasc Disord.* 2017;17(1):1–11.

44. Ziaee M, Mashayekhi S, Ghaffari S, Mahmoudi J, Sarbakhsh P, Garjani A. Predictive Value of Endocan Based on TIMI Risk Score on Major Adverse Cardiovascular Events After Acute Coronary Syndrome. *Angiology.* 2019;70(10):952–9.
45. Silveira DS, Jaeger CP, Hatschbach L, Manenti ERF. Validation of TIMI Risk Score for STEMI. *Int J Cardiovasc Sci.* 2016;29(3):189–97.
46. Nepper-Christensen L, Lønborg J, Høfsten DE, Ahtarovski KA, Bang LE, Helqvist S, et al. Benefit from reperfusion with primary percutaneous coronary intervention beyond 12 hours of symptom duration in patients with ST-segment-elevation myocardial infarction. *Circ Cardiovasc Interv.* 2018;11(9):1–9.
47. Yang HT, Xiu WJ, Zheng YY, Liu F, Gao Y, Ma X, et al. Invasive reperfusion after 12 hours of the symptom onset remains beneficial in patients with ST-segment elevation myocardial infarction: Evidence from a meta-analysis of published data. *Cardiol J.* 2019;26(4):333–42.
48. Khalid U, Jneid H, Denktas AE. The relationship between total ischemic time and mortality in patients with STEMI: Every second counts. *Cardiovasc Diagn Ther.* 2017;7(3):S119–24.
49. Nallamothu BK, Normand SLT, Wang Y, Hofer TP, Brush JE, Messenger JC, et al. Relation between door-to-balloon times and mortality after primary percutaneous coronary intervention over time: A retrospective study. *Lancet.* 2015;385(9973):1114–22.
50. Shahin M, Obeid S, Hamed L, Templin C, Gamperli O, Nietlispach F, et al. Occurrence and Impact of Time Delay to Primary Percutaneous Coronary Intervention in Patients With ST-Segment Elevation Myocardial Infarction. *Cardiol Res.* 2017;8(5):190–8.
51. Hicks KA, Mahaffey KW, Mehran R, Nissen SE, Wiviott SD, Dunn B, et al. 2017 Cardiovascular and Stroke Endpoint Definitions for Clinical Trials. *J Am Coll Cardiol.* 2018;71(9):1021–34.
52. Aleyadeh W, Hutt-Centeno E, Ahmed HM, Shah NP. Hypertension guidelines: Treat patients, not numbers. *Cleve Clin J Med.* 2019;86(3):210–210.
53. Care M. Standards of medical care in diabetes-2018. *Clin diabetes.* 2018;36(1):14–37.
54. Chakraborty S, Majumder B, Sarkar D, Chatterjee S. A Simple Non-invasive ECG Technique to Localize Culprit Vessel Occlusion Site in ST-

Elevation Myocardial Infarction (STEMI) Patients. *J Clin Exp Cardiol.* 2017;8(11).

55. Vicent L, Velásquez-Rodríguez J, Valero-Masa MJ, Díez-Delhoyo F, González-Saldívar H, Bruña V, et al. Predictors of high Killip class after ST segment elevation myocardial infarction in the era of primary reperfusion. *Int J Cardiol.* 2017;248:46–50.
56. Song JX, Zhu L, Lee CY, Ren H, Cao CF, Chen H. Total ischemic time and outcomes for patients with ST-elevation myocardial infarction: Does time of admission make a difference? *J Geriatr Cardiol.* 2016;13(8):658–64.
57. Bucholz EM, Rathore SS, Reid KJ, Jones PG, Chan PS, Rich MW, et al. Body mass index and mortality in acute myocardial infarction patients. *Am J Med.* 2012;125(8):796–803.
58. Zibaeenezhad MJ, Mohammadi SS, Sayadi M, Khorshidi S, Bahramali E, Razeghian-Jahromi I. The impact of diabetes mellitus and hypertension on clinical outcomes in a population of Iranian patients who underwent percutaneous coronary intervention: A retrospective cohort study. *J Clin Hypertens.* 2019;21(11):1647–53.
59. Lolaen C, Rampengan SH, Pangemanan JA. Gambaran Intervensi Koroner Perkutan Primer pada Pasien Infark Miokard Akut dengan Elevasi Segmen ST di RSUP Prof Dr . R . D . J e-Clinic. 2018;6(2):153–61.
60. Afifah HY, Martyarini BS, Yudono DT. Clinical Characteristics of Patients with Acute Myocardial Infarction in Prof. Dr. Margono Soekarjo Regional Public Hospital, Purwokerto. 2020;20(Icch 2019):340–2.
61. Tungsubutra W, Chichareon P, Tresukosol D, Chotinaiwattrakul C, Phankingthongkum R, Wongpraparut N, et al. In-hospital outcomes after rescue percutaneous coronary intervention in patients with ST segment elevation myocardial infarction. *J Med Assoc Thai.* 2016;99(6):645–52.
62. Adhitya AS. Perbandingan Kejadian Kardiovaskular Mayor dengan dan Tanpa Resolusi Depresi Segmen ST Resiprokal pada Penderita Infark Miokard Akut Elevasi Segmen ST yang Mendapatkan Terapi Fibrinolitik Alteplase. Universitas Sumatera Utara; 2018.
63. Laksono BB. Literature Review Efektifitas Terapi Fibrinolitik Dan Ppci (Primary Percutaneous Coronary Intervention) Sebagai Alternative Terapi Revaskularisasi Pada Acute Coronary Syndrome (Acs). *J Kesehat Hesti Wira Sakti.* 2015;3(3):62–8.
64. Reinstadler SJ, Stiermaier T, Eitel C, Saad M, Metzler B, De Waha S, et al. Antecedent hypertension and myocardial injury in patients with reperfused

ST-elevation myocardial infarction. *J Cardiovasc Magn Reson*. 2016;18(1):1–8.

65. Idris H, French JK, Shugman IM, Hopkins AP, Juergens CP, Thomas L. Influence of Age and Gender on Clinical Outcomes Following Percutaneous Coronary Intervention for Acute Coronary Syndromes. *Heart Lung Circ*. 2017;26(6):554–65.
66. Akhtar Z, Abdul Aleem M, Kumar Ghosh P, Monwarul islam AK., Chowdhury F. In-hospital and 30-day major adverse cardiac events in patients referred for ST-segment elevation myocardial infarction in Dhaka , Bangladesh. *Research square*. 2020;1–18.
67. Kanic V, Vollrath M, Naji FH, Sinkovic A. Gender related survival differences in ST-elevation myocardial infarction patients treated with primary PCI. *Int J Med Sci*. 2016;13(6):440–4.
68. Lee MG, Jeong MH, Lee KH, Park KH, Sim DS, Yoon HJ, et al. Prognostic impact of diabetes mellitus and hypertension for mid-term outcome of patients with acute myocardial infarction who underwent percutaneous coronary intervention. *J Cardiol*. 2012;60(4):257–63.
69. Kim YH, Her AY, Jeong MH, Kim BK, Hong SJ, Kim S, et al. Which is the worst risk factor for the long-term clinical outcome? Comparison of long-term clinical outcomes between antecedent hypertension and diabetes mellitus in South Korean acute myocardial infarction patients after stent implantation. *J Diabetes*. 2020;12(2):119–33.
70. Yudha Baharsyah M M. Faktot-faktor yang berhubungan dengan kejadian kardiovaskular mayor selama perawatan pada pasien infark miokard akut dengan elevasi segmen ST yang menjalani intervensi koroner perkutan primer di RSUP DR. M. Djamil Padang. Universitas andalas; 2019.
71. El-Menyar A, Zubaid M, Almahmeed W, Sulaiman K, Alnabti A, Singh R, et al. Killip classification in patients with acute coronary syndrome: Insight from a multicenter registry. *Am J Emerg Med*. 2012;30(1):97–103.
72. Lonborg J, Schoos MM, Kelbæk H, Holmvang L, Steinmetz J, Vejlstrup N, et al. Impact of system delay on infarct size, myocardial salvage index, and left ventricular function in patients with ST-segment elevation myocardial infarction. *Am Heart J*. 2012;164(4):538–46.