

DAFTAR PUSTAKA

1. Wolff K. Disorders of Hair Follicles and Related Disorders. In: Fitzpatrick Color Atlas and Synopsis of Clinical Dermatology. 6th Ed. McGraw-Hill Company; 2009:68-75.
2. Wolff K, LA G, Katz S. Disorders of Hair and Nails. In: Fitzpatrick's Dermatology in General Medicine. 7th Ed. McGraw-Hill Company; 2008:766-9.
3. Legiawati Lili. Alopecia Androgenetik. Dep Ilmu Kesehat Kulit dan Kelamin FK Univ Indonesia /RSUPN dr Cipto Mangunkusumo Jakarta. Published online 2013.
4. G. Severi, R. Sinclair, J.L. Hopper, D.R. English, M.R.E. McCredie, P. Boyle, et al. Androgenetic alopecia in men aged 40–69 years: prevalence and risk factors. 2003;(Br. J. Dermatol):1207–13.
5. Gan DCC SR. Prevalence of male and female pattern hair loss in Maryborough. In: J Investig Dermatol Symp. ; 2005:10.
6. Trüeb RM. Androgenetic alopecia. Eur Handb Dermatological Treat Third Ed. Published online 2015:55-65. doi:10.1007/978-3-662-45139-7_5
7. Alves R GR. Androgenetic Alopecia in Adolescents. In: Oranje AP, Al-Mutairi N ST, ed. Practical Pediatric Dermatology. Controversies in Diagnosis and Treatment. Springer; 2016:187-96.
8. Kim BJ, Kim JY, Eun HC, Kwon OS, Kim MN RB. Androgenetic alopecia in adolescents: a report of 43 cases. J Dermatol. 2006;33(10):696-9.
9. McDonough PH SR. Adolescent androgenic alopecia. Cutis. 2011;88(4):165-8.
10. Hunt N MS. The Psychological impact of alopecia. J Personal Sos Psychol. 2007;20:362-64.
11. Trieu N, Eslick GD. Alopecia and its association with coronary heart disease and cardiovascular risk factors: A meta-analysis. Int J Cardiol. 2014;176(3):687-95. doi:10.1016/j.ijcard.2014.07.079
12. Esen Salman K, Kucukunal NA, Kivanc Altunay I, Aksu Cerman A. Frequency, severity and related factors of androgenetic alopecia in dermatology outpatient clinic: Hospital-based cross-sectional study in Turkey. An Bras Dermatol. 2017;92(1):35-40. doi:10.1590/abd1806-4841.20175241
13. Shaheen A. Clinical Research in Dermatology : Open Access Comprehensive Review of Keloid Formation. Published online 2017.

14. Hirso P, Rajala U, Hiltunen L, Jokelainen J, Keinänen-Kiukaanniemi S, Näyhä S. Obesity and low-grade inflammation among young Finnish men with early-onset alopecia. *Dermatology*. 2007;214(2):125-9. doi:10.1159/000098570
15. Byregowda TS, Kusagur MS, Sugarredy, Muruges. Do High BMI and Severe Androgenetic Alopecia go Hand in Hand ? A Cross-Sectional Study. *Int J Sci Res*. 2016;5(6):2238-40. <https://www.ijsr.net/archive/v5i6/NOV164543.pdf>
16. Su LH, Chen THH. Association of androgenetic alopecia with smoking and its prevalence among Asian men: A community-based survey. *Arch Dermatol*. 2007;143(11):1401-6.
17. Shaikh S, Shaikh S, Shaikh S, Shaikh A, Saleem. S. Prevalence of Hair Loss and Stress As the Cause; a Cross-Sectional Study. *Int J Adv Res*. 2016;4(7):327-33. doi:10.21474/ijar01/924
18. Putri I, Soedibyo S. Tingkat depresi peserta program pendidikan dokter spesialis Ilmu Kesehatan Anak FKUI-RSCM dan faktor-faktor terkait. *Sari Pediatri*. 2011;13(1):1-9.
19. Lolli F, Pallotti F, Rossi A, Fortuna MC, Caro G, Lenzi A, et al. Androgenetic alopecia: a review. *Endocrine*. 2017;57(1):9-17. doi:10.1007/s12020-017-1280-y
20. Messenger, A.G., Sinclair, R.D., Farrant, P. & R. de Berker DA. Acquired Disorders of Hair. In: *Rook's Textbook of Dermatology*.; 2016:2265-342.
21. Habif T. Hair diseases. In: *Clinical Dermatology*. Elsevier Inc.; 2016:923-959.
22. Desmond Chia Chin Gan RS. Androgenetic alopecia. In: *Men's Health*. 3rd ed. ; 2008:353.
23. Hirso P, Rajala U, Hiltunen L, Jokelainen J, Keinänen-Kiukaanniemi S NS. Obesity and low-grade inflammation among young Finnish men with early-onset alopecia. *Dermatology*. 2007;214(125):9.
24. Ahouansou, S., Le Toumelin, P., Crickx, B. & Descamps V. Association of androgenetic alopecia and hypertension. *Eur J Dermatology*. 2007;17(3):220-2.
25. Dardour JC, Pugash E AR. The one-stage preauricular flap for male pattern baldness: Long-term results and risk factors. In: *Plast Reconstr Surg*. ; 1988:907-12.
26. Haley NJ HD. Analysis for nicotine and cotinine in hair to determine cigarette smoker status. *Clin Chem*. 1985;31:1598-600.
27. Agamia NF, Abou Youssif T, El-Hadidy A, El-Abd A. Benign prostatic hyperplasia, metabolic syndrome and androgenic alopecia: Is there a possible relationship? *Arab J Urol*. 2016;14(2):157-162. doi:10.1016/j.aju.2016.01.003
28. Chen W, Kim S, Moon J, Kim H, Kwon D, Won Y. Association of benign prostatic hyperplasia with male pattern baldness. *Urology*. 1998;51:744-8.

29. Chen W, Yang C, Chen G, Wu M, Sheu H, Tzai T. Patients with a large prostate show a higher prevalence of Androgenic Alopecia. *Arch Dermatol*. 2004;296:245-9.
30. Dastgheib L, Shirazi M, Moezzi I, Dehghan S, Sadati MS. Is there a relationship between androgenic alopecia and benign prostatic Hyperplasia ? *Acta Med Iran*. 2015;53(1):30-2.
31. Hadshiew IM, Foitzik K, Arck PC, Paus R. Burden of hair loss: Stress and the underestimated psychosocial impact of telogen effluvium and androgenetic alopecia. *J Invest Dermatol*. 2004;123(3):455-457. doi:10.1111/j.0022-202X.2004.23237.x
32. Putri IA, Soedibyso S. Tingkat Depresi Peserta Program Pendidikan Dokter Spesialis Ilmu Kesehatan Anak FKUI-RSCM dan Faktor-Faktor Terkait. *Sari Pediatri*. 2016;13(1):70. doi:10.14238/sp13.1.2011.70-8
33. Kaliyadan F, Nambiar A, Vijayaraghavan S. Androgenetic alopecia: An update. *Indian J Dermatol Venereol Leprol*. 2013;79(5):613-625. doi:10.4103/0378-6323.116730
34. Blume-peytavi U, Blumeyer A, Tosti A, Finner A, Marvol V, Trakatelli M, et al. S1 guideline for diagnostic evaluation in androgenetic alopecia in men, women and adolescents. *Br J Dermatol*. 2011;164:5-15.
35. Inui S. Trichoscopy for common hair loss diseases: Algorithmic method for diagnosis. *J Dermatol*. 2011;38:71-5.
36. Otberg N, Shapiro J. Hair growth disorders. In: Wolff K, Goldsmith L, Katz S, Gilchrest B, Paller A, Leffell D, eds. *Fitzpatrick's Dermatology in General Medicine*. 8th ed. The McGraw-Hil Companies; 2012:979-1008.
37. Lovinbond. *Depression Anxiety and Stress Scale (DASS)*. 1995;65:1-5.
38. Dahlan, M Sopiudin. Langkah-langkah membuat proposal penelitian bidang kedokteran dan kesehatan. 2nd. Jakarta : CV Sagung Seto, 2016. pp. 80-95.
39. Paik JH, Yoon JB, Sim WY, Kim BS, Kim NI. The prevalence and types of androgenetic alopecia in Korean men and women. *Br J Dermatol* 2001;145:95-8.
40. Brinkman GL, Coates EO Jr. The effect of bronchitis, smoking and occupation on ventilation. *Ann Respir Dis*. 1963;87:684-93.
41. Tang PH, Chia HP, Cheong LL KD. A community study of male androgenetic alopecia in Bishan, Singapore. *Singapore Med J*. 2000;41:202-5.
42. Agamia NF, Abou Youssif T, El-Hadidy A, El-Abd A. Benign prostatic hyperplasia, metabolic syndrome and androgenic alopecia: Is there a possible relationship? *Arab J Urol*. 2016;14(2):157-162. doi:10.1016/j.aju.2016.01.003

43. Arias-Santiago, S., Gutiérrez-Salmerón, M.T., Buendía-Eisman, A., Girón-Prieto, M.S. & Naranjo-Sintes R. Sex hormone-binding globulin and risk of hyperglycemia in patients with androgenetic alopecia. *J Am Acad Dermatol.* 2011;65(1):48-53.
44. Osborn, D. Inheritance of baldness. *J Heredity.* 1916 ; 7, 347-55
45. Ellis JA, Stebbing M, Harrap SB. Polymorphism of the androgen receptor gene is associated with male pattern baldness. *J Investig Dermatol.* 2001; 116: 452-5
46. Tang PH, Chia HP, Cheong LL, Koh D. A community study of male androgenetic alopecia in Bishan, Singapore. *Singapore Med J.* 2000; 41: 202-5
47. Xu F, Sheng YY, Mu ZL, Lou W, Zhou J, et al. Prevalence and types of androgenetic alopecia in Shanghai, China: a community-based study. *Br J Dermatol.* 2009; 160: 629-32
48. Yang, C.C., Hsieh, F.N., Lin, L.Y., Hsu, C.K., Sheu, H.M. & Chen, W. 2014, 'Higher body mass index is associated with greater severity of alopecia in men with male-pattern androgenetic alopecia in Taiwan: A cross-sectional study', *Journal of the American Academy of Dermatology*, vol. 70, no. 2, pp. 297–302.
49. Su, L-H., Chen, T.H-H. 2007, 'Association of Androgenetic Alopecia With Smoking and Its Prevalence Among Asian Men'. *Archives of Dermatology*, Vol. 143, no. 11, pp. 1401-6
50. Rathnayake D, Sinclair R. Male androgenetic alopecia. *Informa J.* 2010; 11(8):1295-304
51. Paul R, Olsen EA, Messenger AG. Hair growth disorders. Dalam: Wolff K, Goldsmith LA, Katz SI, Gilchrist BA, Paller AS, Leffell DJ, penyunting. *Fitzpatrick's dermatology in general medicine.* Edisi ke-7 NewYork: McGraw-Hil Companies, 2008. h. 766-9
52. Nabaie L, Kavand S, Robati RM, Sarrafi-Rad N, Shahgholi L, Meshkat-Razavi G. Androgenic alopecia and insulin resistance: are they really related?. *Clin Exp Dermatol* 2009;34:694-7.
53. Pengsalae N, Tanglertsampan C, Phichawong T, Lee S. Association of early onset androgenetic alopecia and metabolic syndrome in Thai men: a case control study. *J Med Assoc Thai.* 2013;96(8):947-51.
54. Serge A, Philippe LT, Beatrice C, Vincent D. Association of androgenetic alopecia and hypertension. *EJD.* 2007;17:220-2.
55. Hawk E , Brewslow RA , Graubard BI . Male pattern baldness and clinical prostate cancer in the epidemiologic follow-up of the first National Health and Nutrition Examination Survey. *Cancer Epidemiol Biomarkers Prev* 2000 ; 9 : 523 – 7 .

56. Arias-Santiago S, Arrabal-Polo MA, Buendi'a-Eisman A, Arrabal- Marti'n M, Gutie' rrez-Salmero' n MT, Giro' n-Prieto MS, et al. Androgenetic alopecia as an early marker of benign prostatic hyperplasia. *J Am Acad Dermatol* 2012;66:401–8.
57. Hillmer AM, Hanneken S, Ritzmann S, Becker T, Freudenberg J, et al. Genetic variation in the human androgen receptor gene is the major determinant of common early-onset androgenetic alopecia. *Am J Hum Genet.* 2005; 77: 140–8
58. Rajput JR. Role of non androgenic factors in hair loss and hair regrowth. *J Cosmo Trichol.* 2017; 3(2): 118-22
59. Desmond CC, Gan DC, Sinclair RD. Prevalence of male and female pattern hair loss in Maryborough. *J Investig Dermatol Symp Proc.* 2005; 10:184–9
60. Wang TL, Zhou C, Shen XY, Wang XL, Ding S, et al. Prevalence of androgenetic alopecia in china: a community-based study in six cities. *British Journal of Dermatology.* 2010; 162: 843-7.
61. Fahrenkopf AM, Sectish TC, Sharek PJ, Lewin D, Chiang VW, et al. Rates of medication errors among depressed and burnt out residents: prospective cohort study. *BMJ.* epub ahead print
62. West CP, Huschka MM, Novotny PJ, Sloan JA, Kolars JC, Habermann TM, dkk. Association of perceived medical errors with resident distress and empathy. *JAMA* 2006;296:1071-8.
63. Navinés, Ricard & Martín-Santos, Rocío & Olivé, Victoria & Valdés, Manuel. (2016). Work-related stress: Implications for physical and mental health. *Medicina Clínica (English Edition).* 146. 10.1016/j.medcle.2016.06.010.
64. Legiawati L. Jenis kerontokan rambut dan kebutakan pasien poliklinik kulit dan kelamin RSUPN. Dr. Cipto Mangunkusumo Tahun 2009- 2011. *MDVI* 2013;40:159-63.

