

DAFTAR PUSTAKA

- Agarwal, S. and Krishnamurty, K. (2019) *Histology Skin*, StatPearls Publishing.
- Agrawal, O. P. and Agrawal, S. (2012) 'An Overview of New Drug Delivery System : Microemulsion quaternary phase diagrams can describe the phase', *Asian Journal of Pharmaceutical Science & Technology*, 2(1), pp. 5–12.
- Aldag, C., Teixeira, D. N. and Leventhal, P. S. (2016) 'Skin rejuvenation using cosmetic products containing growth factors , cytokines , and matrikines : a review of the literature', pp. 411–419.
- Ángeles, T. D. L. *et al.* (2018) 'Sensory analysis of cosmetic formulations made with essential oils of *Aristeguietia glutinosa* (matico) and *Ocotea quixos* (ishpingo)'. doi: 10.15171/ijpni.2018.05.
- Anonim (1979) *Farmakope Indonesia*. Edisi III. Jakarta: Departemen Kesehatan Republik Indonesia.
- Anonim (1995) *Farmakope Indonesia*. Edisi IV. Jakarta: Departemen Kesehatan Republik Indonesia.
- Ata, S. T. W. *et al.* (2016) 'Isolasi Kolagen Dari Kulit Dan Tulang Ikan Cakalang (*Katsuwonus pelamis*)', *Journal of Pharmaceutical and Medicinal Science*, 1(1), pp. 27–30.
- ATCC (2014) *Animal Cell Culture Guide*. Manassas.
- Badan Standardisasi Nasional (1996) 'Sediaan Tabir Surya', *Dewan Standardisasi Nasional*, 16(4399), pp. 1–3.
- Barel, A. O., Paye, M. and Howard, I. M. (2009) *Handbook of Cosmetic Science and Technology*. Third Edit. Edited by I. Healthcare. New York.
- Berneburg, M. *et al.* (1997) 'Chronically Ultraviolet-exposed Human Skin Shows a Higher Mutation Frequency of Mitochondrial DNA as Compared to Unexposed Skin and the Hematopoietic System', *Photochemistry and Photobiology*, 66(2), pp. 271–275. doi: 10.1111/j.1751-1097.1997.tb08654.x.
- Berneburg, M., Plettenberg, H. and Krutmann, J. (2000) 'Photoaging of human skin', pp. 239–244.
- Bos, J. D. and Meinardi, M. M. H. M. (2000) 'The 500 Dalton rule for the skin penetration of chemical compounds and drugs', *Experimental Dermatology*, 9(3), pp. 165–169. doi: 10.1034/j.1600-0625.2000.009003165.x.

- BPOM RI (2019) *Peraturan Badan Pengawas Obat dan Makanan Nomor 23 Tahun 2019 Tentang Persyaratan Teknis Bahan Kosmetik, Bpom Ri.*
- Brandt, F. S., Cazzaniga, A. and Hann, M. (2011) 'Cosmeceuticals: Current Trends and Market Analysis', *Seminars in Cutaneous Medicine and Surgery*, 30(3), pp. 141–143. doi: 10.1016/j.sder.2011.05.006.
- Caroll, M. (2020) *10 Best Stem Cell Beauty Products On The Market Today*, *Bioinformant.com*. Available at: <https://bioinformant.com/beauty-products-stem-cells/>.
- Cohen, M. and Cohen, J. (2010) 'Umbilical Cord Stem Cell Secreted Product Derived Topical Compositions And Methods Of Use Thereof'. Available at: <https://patentimages.storage.googleapis.com/3b/c9/82/c283c7b24afe69/US20100019677A1.pdf>.
- Das, A. and Ahmed, A. B. (2017) 'Formulation and evaluation of transdermal patch of indomethacin containing patchouli oil as natural penetration enhancer', *Asian Journal of Pharmaceutical and Clinical Research*, 10(11), pp. 320–325. doi: 10.22159/ajpcr.2017.v10i11.20926.
- Dewi, R., Anwar, E. and Yunita, K. S. (2014) 'Uji Stabilitas Fisik Formula Krim yang Mengandung Ekstrak Kacang Kedelai (Glycine max) Abstrak', 1(3), pp. 194–208.
- Dominici, M. *et al.* (2006) 'Minimal criteria for defining multipotent mesenchymal stromal cells. The International Society for Cellular Therapy position statement', *Cytotherapy*, 8(4), pp. 315–317. doi: 10.1080/14653240600855905.
- Dominik, I. *et al.* (2015) 'Activation of TGF- β : a gateway to skin rejuvenation in skin to unlock TGF- β potential', 10(December), pp. 6–11.
- Draelos, Z. D. (2011) *Cosmetic Dermatology: Products and Procedures*. USA: Wiley-Blackwell.
- El-Domyati, M. *et al.* (2020) 'Facial rejuvenation using stem cell conditioned media combined with skin needling: A split face comparative study', *Journal of Cosmetic Dermatology*, 19(9), pp. 2404–2410. doi: 10.1111/jocd.13594.
- Elmitra (2018) *Dasar-Dasar Farmasetika dan Sediaan Semi Solid*. Yogyakarta: Deepublish.
- Eurell, J. A. C., Frappier, B. L. and Dellmann, H.-D. (2006) *Dellmann's textbook of veterinary histology*. Iowa: Blackwell Pub.
- Flament, F. *et al.* (2015) 'Facial skin pores: A multiethnic study', *Clinical, Cosmetic and Investigational Dermatology*, 8, pp. 85–93. doi: 10.2147/CCID.S74401.

- Foldvari, M. (2000) 'Non-invasive administration of drugs through the skin: Challenges in delivery system design', *Pharmaceutical Science and Technology Today*, 3(12), pp. 417–425. doi: 10.1016/S1461-5347(00)00317-5.
- Fox, L. T. *et al.* (2011) 'Transdermal drug delivery enhancement by compounds of natural origin', *Molecules*, 16(12), pp. 10507–10540. doi: 10.3390/molecules161210507.
- Fu, X. *et al.* (2007) 'Adipose tissue extract enhances skin wound healing', *Wound Repair and Regeneration*, 15(4), pp. 540–548. doi: 10.1111/j.1524-475X.2007.00262.x.
- Hagen, Ø., Solberg, C. and Johnston, I. A. (2008) 'Activity of aspartate (Cathepsin D), cysteine proteases (Cathepsins B, B + L, and H), and matrix metalloproteinase (Collagenase) and their influence on protein and water-holding capacity of muscle in commercially farmed atlantic halibut (*Hippoglossus hippo*)', *Journal of Agricultural and Food Chemistry*, 56(14), pp. 5953–5959. doi: 10.1021/jf801215b.
- Halim, D. *et al.* (2010) *Stem Cell: Dasar Teori & Aplikasi Klinis*. Erlangga.
- Hallstar (2002) 'Olivem® 1000 The First Emulsifying Active Ingredient', *Product Literature*. Chicago: The Hallstar Company, pp. 1–10.
- Hartono, B. (2016) 'Sel Punca : Karakteristik , Potensi dan Aplikasinya', *J. Kedokteran Meditek*, 22(60), pp. 72–75.
- Herman, A. and Herman, A. P. (2015) 'Essential oils and their constituents as skin penetration enhancer for transdermal drug delivery: A review', *Journal of Pharmacy and Pharmacology*, 67(4), pp. 473–485. doi: 10.1111/jphp.12334.
- Invitrogen (2009) 'DMEM Dulbecco's Modified Eagle Media'.
- Jayaraman, P. *et al.* (2013) 'Stem cells conditioned medium: A new approach to skin wound healing management', *Cell Biology International*, 37(10), pp. 1122–1128. doi: 10.1002/cbin.10138.
- Kalangi, S. J. R. (2014) 'Histofisiologi Kulit', *Jurnal Biomedik (Jbm)*, 5(3), pp. 12–20. doi: 10.35790/jbm.5.3.2013.4344.
- Kale, S. N. and Deore, S. L. (2016) 'Emulsion micro emulsion and nano emulsion: A review', *Systematic Reviews in Pharmacy*, 8(1), pp. 39–47. doi: 10.5530/srp.2017.1.8.
- Kim, H. J. *et al.* (2020) 'A study on clinical effectiveness of cosmetics containing human stem cell conditioned media', *Biomedical Dermatology*, 4(1), pp. 1–11. doi: 10.1186/s41702-020-0056-9.
- Kim, J. *et al.* (2011) 'Inhibition effect of *Gynura procumbens* extract on UV-B-induced matrix-metalloproteinase expression in human dermal

fibroblasts', *Journal of Ethnopharmacology*, 137(1), pp. 427–433. doi: 10.1016/j.jep.2011.04.072.

- Kim, K. H. *et al.* (2020) 'The effect of three-dimensional cultured adipose tissue-derived mesenchymal stem cell-conditioned medium and the antiaging effect of cosmetic products containing the medium', *Biomedical Dermatology*, 4(1), pp. 1–12. doi: 10.1186/s41702-019-0053-z.
- Kim, S. K. *et al.* (2002) 'Purification and characterization of a collagenolytic protease from the filefish, *Novodon modestrus*', *Journal of Biochemistry and Molecular Biology*, 35(2), pp. 165–171. doi: 10.5483/bmbrep.2002.35.2.165.
- Kim, W. S. *et al.* (2007) 'Wound healing effect of adipose-derived stem cells: A critical role of secretory factors on human dermal fibroblasts', *Journal of Dermatological Science*, 48(1), pp. 15–24. doi: 10.1016/j.jdermsci.2007.05.018.
- Kwon, T. R. *et al.* (2016) 'Conditioned medium from human bone marrow-derived mesenchymal stem cells promotes skin moisturization and effacement of wrinkles in UVB-irradiated SKH-1 hairless mice', *Photodermatology Photoimmunology and Photomedicine*, 32(3), pp. 120–128. doi: 10.1111/phpp.12224.
- Lawrence, M. J. and Rees, G. D. (2012) 'Microemulsion-based media as novel drug delivery systems', *Advanced Drug Delivery Reviews*, 64(SUPPL.), pp. 175–193. doi: 10.1016/j.addr.2012.09.018.
- Lephart, E. D. (2016) 'Skin Aging and Oxidative Stress: Equol's Anti-Aging Effects via Biochemical and Molecular Mechanisms', *Ageing Research Reviews*. doi: 10.1016/j.arr.2016.08.001.
- Lestari, N. K. L., Suardana, I. W. and Sukrama, I. D. M. (2019) 'Karakteristik Fisikokimia dan Uji Aktivitas Antimikroba Bakteriosin dari Isolat Bakteri Asam Laktat 15B hasil Isolasi Kolon Sapi Bali', *Buletin Veteriner Udayana*, (21), p. 65. doi: 10.24843/bulvet.2019.v11.i01.p11.
- Li, M. *et al.* (2017) 'Umbilical cord-derived mesenchymal stromal cell-conditioned medium exerts in vitro antiaging effects in human fibroblasts', *Cytotherapy*, 19(3), pp. 371–383. doi: 10.1016/j.jcyt.2016.12.001.
- Liu, H. *et al.* (2012) 'Original Article Anti-oxidant pathways are stimulated by mesenchymal stromal cells in renal repair after ischemic injury', *Cytotherapy*, 14(2), pp. 162–172. doi: 10.3109/14653249.2011.613927.
- Liu, Q. *et al.* (2013) 'Conditioned serum-free medium from umbilical cord mesenchymal stem cells has anti-photoaging properties',

Biotechnology Letters, 35(10), pp. 1707–1714. doi: 10.1007/s10529-013-1242-2.

- Liu, Z. *et al.* (2017) ‘Potential of bone marrow mesenchymal stem cells in rejuvenation of the aged skin of rats’, *Biomedical Reports*, 6(3), pp. 279–284. doi: 10.3892/br.2017.842.
- Ma’at, S. (2011) *Teknik Dasar Kultur Sel*. Yogyakarta: Airlangga University Press.
- Mardhiani, Y. D. *et al.* (2018) ‘Formulasi dan Stabilitas Sediaan Serum dari Ekstrak Kopi Hijau (*Coffea canephora* var. *Robusta*) sebagai Antioksidan’, *Indonesia Natural Research Pharmaceutical Journal Universitas 17 Agustus 1945 Jakarta*, 2(2), pp. 19–33.
- Matangi, S. P. *et al.* (2014) ‘Formulation and evaluation of anti aging poly herbal cream’, *International Journal of Pharmaceutical Sciences Review and Research*, 24(2), pp. 133–136.
- Mather, J. P. and Roberts, P. E. (1998) *Introduction to Cell and Tissue Culture Theory and Technique*. Edited by J. P. Mather and P. E. Roberts. New York: Plenum Press.
- Miani, J. (2018) *Apa Itu Essence? Kenapa Skincare Ini Penting Untuk Menjaga Kecantikan Kulit? Cari Tahu di Sini!*, *The Shonet*. Available at: theshonet.com/articles/apa-itu-essence-kenapa-skincare-ini-penting-untuk-menjaga-kecantikan-kulit-cari-tahu-di-sini.
- Mitsui, T. (1998) *New Cosmetic Science*. Amsterdam: Elsevier Science B.V.
- Osugi, M. *et al.* (2012) ‘Conditioned media from mesenchymal stem cells enhanced bone regeneration in rat calvarial bone defects’, *Tissue Engineering - Part A*, 18(13–14), pp. 1479–1489. doi: 10.1089/ten.tea.2011.0325.
- Park, B. S. *et al.* (2008) ‘Adipose-derived stem cells and their secretory factors as a promising therapy for skin aging’, *Dermatologic Surgery*, 34(10), pp. 1323–1326. doi: 10.1111/j.1524-4725.2008.34283.x.
- Park, B. S. *et al.* (2010) ‘Hair growth stimulated by conditioned medium of adipose-derived stem cells is enhanced by hypoxia: Evidence of increased growth factor secretion’, *Biomedical Research*, 31(1), pp. 27–34. doi: 10.2220/biomedres.31.27.
- Patel, G. K. *et al.* (2019) ‘Comparative analysis of exosome isolation methods using culture supernatant for optimum yield, purity and downstream applications’, *Scientific Reports*, 9(1), pp. 1–10. doi: 10.1038/s41598-019-41800-2.
- Pawitan, J. A. (2014) ‘Prospect of Stem Cell Conditioned Medium in’, *BioMed Research International*, 2014, pp. 7–9. doi: 10.1155/2014/965849.

- Prafita, N., Lucida, H. and Fitriani, L. (2019) *Optimasi Formula Krim Tabir Surya Ekstrak Tomat (Solanum lycopersicum L.) Terpurifikasi Menggunakan Surfaktan Olivem dan Fase Minyak VCO*. Universitas Andalas.
- Priani, S. E. *et al.* (2013) 'Formulasi Sediaan Emulgel Untuk Penghantaran Transdermal Ketoprofen', XXXVIII(1), pp. 37–42.
- Rabe, J. H. *et al.* (2006) 'Photoaging: Mechanisms and repair', pp. 1–19. doi: 10.1016/j.jaad.2005.05.010.
- Ramos-e-Silva, M. *et al.* (2013) 'Anti-aging cosmetics: Facts and controversies', *Clinics in Dermatology*, 31(6), pp. 750–758. doi: 10.1016/j.clindermatol.2013.05.013.
- Rifati, L. *et al.* (2012) *Laporan Penelitian: Induksi In-Vitro Sel Punca Mesenkim dari Tali Pusat Manusia Menjadi Sel Punca Limbal*. Jakarta.
- Rizal, R. *et al.* (2019) 'Effect of Serial Passage on Growth Kinetics, Biological Properties, and Differentiation Into Adipocytes of Human Whartonâ€™S Jelly-Derived Mesenchymal Stem Cells', *Majalah Kedokteran Bandung*, 51(3), pp. 127–133. doi: 10.15395/mkb.v51n3.1767.
- Rowe, R. C., Sheskey, P. J. and Quinn, M. E. (2006) *Handbook of Pharmaceutical Excipients*. 6th edn. USA: Pharmaceutical Press and American Pharmacists Association.
- Sasaki, M. *et al.* (2008) 'Mesenchymal Stem Cells Are Recruited into Wounded Skin and Contribute to Wound Repair by Transdifferentiation into Multiple Skin Cell Type', *The Journal of Immunology*, 180(4), pp. 2581–2587. doi: 10.4049/jimmunol.180.4.2581.
- Sasidharan, S., Joseph, P. and Junise (2014) 'Formulation and Evaluation of Fairness Serum Using Polyherbal Extracts', *International Journal of Pharmacy*, 4(3), pp. 105–112. Available at: <http://www.pharmascholars.com>.
- Sato, C. *et al.* (2018) 'Conditioned medium obtained from amnion-derived mesenchymal stem cell culture prevents activation of keloid fibroblasts', *Plastic and Reconstructive Surgery*, 141(2), pp. 390–398. doi: 10.1097/PRS.0000000000004068.
- Sekar, M., Sivalingam, P. and Mahmud, A. (2017) 'Formulation and Evaluation of Novel Antiaging Cream Containing', *International Journal of Pharmaceutical Sciences and Research*, 8(3), pp. 1056–1065. doi: 10.13040/IJPSR.0975-8232.8(3).1056-65.
- Seo, K. Y. *et al.* (2012) 'Skin rejuvenation by microneedle fractional radiofrequency treatment in Asian skin; Clinical and histological analysis', *Lasers in Surgery and Medicine*, 44(8), pp. 631–636. doi:

10.1002/ism.22071.

- Sharma, B., Sharma, A. and Arora, S. (2012) 'Formulation , Optimization and Evaluation of Atorvastatin Calcium Loaded Microemulsion', pp. 1–7.
- Sigma-Aldrich (2016) 'Collagenase Activity Colorimetric Assay', *Technical bulletin*. USA, pp. 2–4. Available at: <https://www.sigmaaldrich.com/content/dam/sigmaaldrich/docs/Sigma/Bulletin/2/mak293bul.pdf>.
- Simper, D. *et al.* (2010) 'Comparative proteomics profiling reveals role of smooth muscle progenitors in extracellular matrix production', *Arteriosclerosis, Thrombosis, and Vascular Biology*, 30(7), pp. 1325–1332. doi: 10.1161/ATVBAHA.110.204651.
- Sinko, P. J. (2011) *Martin Farmasi Fisika dan Ilmu Farmasetika*. Edisi 5. Jakarta: Penerbit Buku Kedokteran EGC.
- Siswanto, F. M. *et al.* (2014) 'The Balancing of Physical Fitness Increase The Activity of Endogenous Stem Cell for Antiaging', 2(1), pp. 1–9.
- Sohn, S. J. *et al.* (2018) 'Anti-aging Properties of Conditioned Media of Epidermal Progenitor Cells Derived from Mesenchymal Stem Cells', *Dermatology and Therapy*, 8(2), pp. 229–244. doi: 10.1007/s13555-018-0229-2.
- Stefanini, M. O. *et al.* (2008) 'A compartment model of VEGF distribution in blood, healthy and diseased tissues', *BMC Systems Biology*, 2, pp. 1–25. doi: 10.1186/1752-0509-2-77.
- Swastika, A. and Purwanto, M. (2015) 'Antioxidant Activity of Cream Dosage Form of Tomato Extract (*Solanum lycopersicum* L.)', *Traditional Medicine Journal*, 18(3), pp. 132–140. doi: 10.22146/tradmedj.8214.
- Syamsuni (2006) *Ilmu Resep*. Jakarta: EGC.
- Tarwoto, Wartonah and Aryani, R. (2009) *Anatomi dan fisiologi: untuk mahasiswa keperawatan*. Jakarta: Trans Info Media.
- Task Force Committee for Evaluation of Anti-Aging Function (2007) 'Guidelines for evaluation of anti-wrinkle products (Guidelines for evaluation of cosmetic functions)', *Journal of Japanese Cosmetic Science Society*, 31(4), pp. 411–431.
- Thomas, B. J. and Finnin, B. C. (2004) 'The transdermal revolution', *Drug Discovery Today*, 9(16), pp. 697–703. doi: 10.1016/S1359-6446(04)03180-0.
- Tiwari, A. (2017) *Practical Biochemistry: A Student Companion*. Guwahati: Lambert Academy Publishing.
- Tobin, D. J. (2017) 'Introduction to skin aging', *Journal of Tissue Viability*, 26(1),

pp. 37–46. doi: 10.1016/j.jtv.2016.03.002.

- Tranggano, R. I. and Latifah, F. (2014) *Buku Pegangan Dasar Kosmetologi*. Edited by J. Djajadisastra and P. Adimukti. Jakarta: Sagung Seto.
- Valacchi, G. *et al.* (2003) ‘Induction of stress proteins and MMP-9 by 0.8 ppm of ozone in murine skin’, *Biochemical and Biophysical Research Communications*, 305(3), pp. 741–746. doi: 10.1016/S0006-291X(03)00812-X.
- Vierkötter, A. and Krutmann, J. (2012) ‘Environmental influences on skin aging and ethnic-specific manifestations’, *Dermato-Endocrinology*, 4(3), pp. 227–231. doi: 10.4161/derm.19858.
- Voigt, R. (1994) *Buku Pelajaran Teknologi Farmasi*. Yogyakarta: Gadjah Mada University Press.
- Wade, A. and Weller, P. J. (1994) *Handbook of pharmaceutical excipients*. 2nd edn. Washington: American Pharmaceutical Association.
- Walker, R. B. and Smith, E. W. (1996) ‘The role of percutaneous penetration enhancers’, *Advanced Drug Delivery Reviews*, 18(3), pp. 295–301. doi: 10.1016/0169-409X(95)00078-L.
- Walter, M. N. M. *et al.* (2010) ‘Mesenchymal stem cell-conditioned medium accelerates skin wound healing: An in vitro study of fibroblast and keratinocyte scratch assays’, *Experimental Cell Research*, 316(7), pp. 1271–1281. doi: 10.1016/j.yexcr.2010.02.026.
- Wasitaatmadja, S. M. (1997) *Penuntun Ilmu Kosmetik Medik*. Edited by U. Press. Jakarta.
- Widowati, W. *et al.* (2015) ‘Conditioned medium from normoxia (WJMSCs-norCM) and hypoxia-treated WJMSCs (WJMSCs-hypoCM) in inhibiting cancer cell proliferation’, *Biomarkers and Genomic Medicine*, 7(1), pp. 8–17. doi: 10.1016/j.bgm.2014.08.008.
- Widowati, W. *et al.* (2016) ‘Antioxidant and anti aging assays of *Oryza sativa* extracts, vanillin and coumaric acid’, *Journal of Natural Remedies*, 16(3), pp. 88–99. doi: 10.18311/jnr/2016/7220.
- Xu, Yang *et al.* (2015) ‘Antiphotoaging Effect of Conditioned Medium of Dedifferentiated Adipocytes on Skin in Vivo and in Vitro: A Mechanistic Study’, *Stem Cells and Development*, 24(9), pp. 1096–1111. doi: 10.1089/scd.2014.0321.
- Yasui, H. and Sakurai, H. (2003) ‘Age-dependent generation of reactive oxygen species in the skin of live hairless rats exposed to UVA light’, *Experimental Dermatology*, 12(5), pp. 655–661. doi: 10.1034/j.1600-0625.2003.00033.x.
- Yin, L., Morita, A. and Tsuji, T. (2000) ‘Alterations of extracellular matrix

induced by tobacco smoke extract’, *Archives of Dermatological Research*, 292(4), pp. 188–194. doi: 10.1007/s004030050476.

Yin, L., Morita, A. and Tsuji, T. (2001) ‘Skin aging induced by ultraviolet exposure and tobacco smoking: Evidence from epidemiological and molecular studies’, *Photodermatology Photoimmunology and Photomedicine*, 17(4), pp. 178–183. doi: 10.1034/j.1600-0781.2001.170407.x.

Zara, S. *et al.* (2013) ‘Development and Aging Are Oxygen-Dependent and Correlate with VEGF and NOS along Life Span’, pp. 223–228. doi: 10.1007/978-94-007-4549-0.

Zhang, S. and Duan, E. (2018) ‘Fighting against Skin Aging: The Way from Bench to Bedside’, *Cell Transplantation*, 27(5), pp. 729–738. doi: 10.1177/0963689717725755.

Zhang, W. and Liu, L. (2013) ‘Study on the Formation and Properties of Liquid Crystal Emulsion in Cosmetic’, 2013(June), pp. 139–144.

