

DAFTAR PUSTAKA

1. Juzar D DS, Tobing D, Firdaus I, Widyantoro B, Rossimarina V, et al. Pedoman tata laksana Sindrom Koroner Akut. 2018. Edisi IV. Jakarta. PERKI;2018:1-2
2. Bi X, Yang C, Song Y, Yuan J, Cui J, Hu F, et al. Quantitative fragmented QRS has a good diagnostic value on myocardial fibrosis in hypertrophic obstructive cardiomyopathy based on clinical-pathological study. *BMC cardiovascular disorders.* 2020;20(1):1-9.
3. Das MK, Suradi H, Maskoun W, Michael MA, Shen C, Peng J, et al. Fragmented wide QRS on a 12-lead ECG: a sign of myocardial scar and poor prognosis. *Circulation: Arrhythmia and Electrophysiology.* 2008;1(4):258-68.
4. Das MK, Khan B, Jacob S, Kumar A, Mahenthiran J. Significance of a fragmented QRS complex versus a Q wave in patients with coronary artery disease. *Circulation.* 2006;113(21):2495-501.
5. Das MK, Michael MA, Suradi H, Peng J, Sinha A, Shen C, et al. Usefulness of fragmented QRS on a 12-lead electrocardiogram in acute coronary syndrome for predicting mortality. *The American journal of cardiology.* 2009;104(12):1631-7.
6. Pietrasik G, Zaręba W. QRS fragmentation: diagnostic and prognostic significance. *Cardiology journal.* 2012;19(2):114-21.
7. Mahfouz RA, Arab M, Abdelhamid M, Elzayat A. Fragmented QRS complex is an independent predictor of plaque burden in patients at intermediate risk of coronary artery disease. *Indian heart journal.* 2019;71(5):394-9.
8. Take Y, Morita H. Fragmented QRS: what is the meaning? *Indian pacing and electrophysiology journal.* 2012;12(5):213-25.
9. Rahman HA, Ghany MA, Youssef AAA. Correlation of fragmented QRS complexes with the severity of CAD (using Syntax score) in patients with non-ST elevation acute coronary syndromes. *The Egyptian Heart Journal.* 2016;68(2):125-9.
10. Cetin M, Kocaman SA, Canga A, Durakoglugil ME, Erdogan T, Satiroglu O, et al. The independent relationship between systemic inflammation and fragmented QRS complexes in patients with stable angina pectoris. *Kardiologia Polska (Polish Heart Journal).* 2012;70(7):668-75.
11. Kocaman SA, Çetin M, Kiriş T, Erdoğan T, Çanga A, Durakoğlugil E, et al. The importance of fragmented QRS complexes in prediction of myocardial infarction and reperfusion parameters in patients undergoing primary percutaneous coronary intervention. *Turk Kardiyol Dern Ars.* 2012;40(3):213-22.
12. Korkmaz A, Yıldız A, Demir M, Ozyazgan B, Sahan E, Acar B, et al. The relationship between fragmented QRS and functional significance of coronary lesions. *Journal of electrocardiology.* 2017;50(3):282-6.
13. Gensini GG. A more meaningful scoring system for determining the severity of coronary heart disease. *Am J cardiol.* 1983;51:606.
14. Collet J-P, Thiele H, Barbato E, Barthélémy O, Bauersachs J, Bhatt DL, et al. 2020 ESC Guidelines for the management of acute coronary syndromes in

- patients presenting without persistent ST-segment elevationThe Task Force for the management of acute coronary syndromes in patients presenting without persistent ST-segment elevation of the European Society of Cardiology (ESC). *European heart journal*. 2020;8-10
- 15. McManus DD, Gore J, Yarzebski J, Spencer F, Lessard D, Goldberg RJ. Recent trends in the incidence, treatment, and outcomes of patients with STEMI and NSTEMI. *The American journal of medicine*. 2011;124(1):40-7.
 - 16. Yuniadi Y HD, Siswanto BB, Santoso A, Zamroni D, et al Buku Ajar Kardiovaskular. Jakarta. 2017. 177-86 p.
 - 17. Kumar A, Cannon CP, editors. Acute coronary syndromes: diagnosis and management, part I. Mayo Clinic Proceedings; 2009: Elsevier.
 - 18. Zipes DP, Libby P, Bonow RO, Mann DL, Tomaselli GF. Braunwald's heart disease e-book: A textbook of cardiovascular medicine: Elsevier Health Sciences; 2019; 2673-2776p
 - 19. Kloner RA. Stunned and hibernating myocardium: where are we nearly 4 decades later? *Journal of the American Heart Association*. 2020;9(3):e015502.
 - 20. Lilly LS. Pathophysiology of heart disease: a collaborative project of medical students and faculty: Lippincott Williams & Wilkins; 2012;75-112
 - 21. Neglia D, Rovai D, Caselli C, Pietila M, Teresinska A, Aguadé-Bruix S, et al. Detection of significant coronary artery disease by noninvasive anatomical and functional imaging. *Circulation: Cardiovascular Imaging*. 2015;8(3):e002179.
 - 22. Issa Z, Miller JM, Zipes DP. Clinical Arrhythmology and Electrophysiology: A Companion to Braunwald's Heart Disease E-Book: Expert Consult: Online and Print: Elsevier Health Sciences; 2019;1-14
 - 23. Brohet C. Fragmentation of the QRS complex: the latest electrocardiographic craze? : Taylor & Francis; 2019;3
 - 24. Mittal SR. Fragmented QRS: A simple electrocardiographic prognostic marker in cardiovascular disease. *Journal of clinical and preventive cardiology*. 2016;5(3):94.
 - 25. Jain R, Singh R, Yamini S, K Das M. Fragmented ECG as a risk marker in cardiovascular diseases. *Current Cardiology Reviews*. 2014;10(3):277-86.
 - 26. Morita H, Kusano KF, Miura D, Nagase S, Nakamura K, Morita ST, et al. Fragmented QRS as a marker of conduction abnormality and a predictor of prognosis of Brugada syndrome. *Circulation*. 2008;118(17):1697.
 - 27. Cho H, Yoon J, Kim N, Jang S, Bae M, Lee J, et al. Predictive value of a fragmented QRS complex in diagnosing patients with myocardial ischemia. *Clinical cardiology*. 2019;42(3):379-84.
 - 28. Ferdous MAR, Mandal M, Kabir FI, Amin MR, Ahsan SA, Mahbub SEE, et al. Angiographic Severity of Coronary Artery Disease among Patients with Non-ST Elevated Acute Coronary Syndrome having Fragmented-QRS in ECG. *University Heart Journal*. 2020;16(1):28-32.
 - 29. France R, Formolo J, Penney D. Value of notching and slurring of the resting QRS complex in the detection of ischemic heart disease. *Clinical cardiology*. 1990;13(3):190-6.

30. Neeland IJ, Patel RS, Eshtehardi P, Dhawan S, McDaniel MC, Rab ST, et al. Coronary angiographic scoring systems: an evaluation of their equivalence and validity. *American heart journal*. 2012;164(4):547-52. e1.
31. Huang G, Zhao J-l, Du H, Lan X-b, Yin Y-h. Coronary score adds prognostic information for patients with acute coronary syndrome. *Circulation Journal*. 2010;0912250569-.
32. Sinning C, Lillpopp L, Appelbaum S, Ojeda F, Zeller T, Schnabel R, et al. Angiographic score assessment improves cardiovascular risk prediction: the clinical value of SYNTAX and Gensini application. *Clinical Research in Cardiology*. 2013;102(7):495-503.
33. Yildirim E, Iyisoy A, Celik M, Yuksel UC, Acikel C, Bugan B, et al. The Relationship Between Gensini Score and In-Hospital Mortality in Patients with ST-Segment Elevation Myocardial Infarction. *International Journal of Cardiovascular Sciences*. 2017;30(1):32-41.
34. Pattarabanjird T, Cress C, Nguyen A, Taylor A, Bekiranov S, McNamara C. A Machine Learning Model Utilizing a Novel SNP Shows Enhanced Prediction of Coronary Artery Disease Severity. *Genes*. 2020;11(12):1446.
35. Murakami N, Kokubu N, Nishida J, Hase M, Fujito T, Kawamukai M, et al. do two scoring systems of coronary stenosis, syntax score and Gensini score, similarly predict clinical outcome after Pci in patients with stable angina pectoris? *Circulation*. 2014;130(suppl_2):A15183-A.
36. Li X-T, Fang H, Li D, Xu F-Q, Yang B, Zhang R, et al. Association of platelet to lymphocyte ratio with in-hospital major adverse cardiovascular events and the severity of coronary artery disease assessed by the Gensini score in patients with acute myocardial infarction. *Chinese medical journal*. 2020;133(4):415.
37. Qin W, Yang Y, Li X, Men L, Guo J, Liu F, et al. Value of mean platelet volume and Gensini score on predicting short-term outcome in acute ST segment elevation myocardial infarction patient post emergency percutaneous coronary intervention. *Zhonghua xin xue guan bing za zhi*. 2015;43(1):22-5.
38. Wright RS, Anderson JL, Adams CD, Bridges CR, Casey DE, Ettinger SM, et al. 2011 ACCF/AHA focused update incorporated into the ACC/AHA 2007 guidelines for the management of patients with unstable angina/non-ST-elevation myocardial infarction: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. *Journal of the American College of Cardiology*. 2011;57(19):e215-e367.
39. Anderson JL, Adams CD, Antman EM, Bridges CR, Califf RM, Casey DE, Jr., et al. ACC/AHA 2007 guidelines for the management of patients with unstable angina/non-ST-Elevation myocardial infarction: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Writing Committee to Revise the 2002 Guidelines for the Management of Patients With Unstable Angina/Non-ST-Elevation Myocardial Infarction) developed in collaboration with the American College of Emergency Physicians, the Society for Cardiovascular Angiography and Interventions, and the Society of Thoracic Surgeons endorsed by the American Association of Cardiovascular and Pulmonary Rehabilitation and

- the Society for Academic Emergency Medicine. *Journal of the American College of Cardiology*. 2007;50(7):e1-e157. Epub 2007/08/19.
40. Piepoli MF, Hoes AW, Agewall S, Albus C, Brotons C, Catapano AL, et al. Guidelines: Editor's choice: 2016 European Guidelines on cardiovascular disease prevention in clinical practice: The Sixth Joint Task Force of the European Society of Cardiology and Other Societies on Cardiovascular Disease Prevention in Clinical Practice (constituted by representatives of 10 societies and by invited experts) Developed with the special contribution of the European Association for Cardiovascular Prevention & Rehabilitation (EACPR). *European heart journal*. 2016;37(29):2315.
41. De Lemos J, Omland T. Chronic coronary artery disease: a companion to Braunwald's heart disease E-book: elsevier Health Sciences; 2018;3;194
42. Kang MJ, Oh Y-M, Lee JC, Kim DG, Park MJ, Lee MG, et al. Lung matrix metalloproteinase-9 correlates with cigarette smoking and obstruction of airflow. *Journal of Korean medical science*. 2003;18(6):821.
43. Soelistijo SA, Novida H, Rudijanto A, Soewondo P, Suastika K, Manaf A, et al. Definisi, Patogenesis, Klasifikasi Diabetes Melitus Tipe 2. In: Rudijanto A, editor. *Konsensus Pengelolaan dan Pencegahan Diabetes Melitus Tipe 2 di Indonesia*. Jakarta: PERKENI; 2015: 6-11.
44. Guo R, Li Y, Xu Y, Tang K, Li W. Significance of fragmented QRS complexes for identifying culprit lesions in patients with non-ST-elevation myocardial infarction: a single-center, retrospective analysis of 183 cases. *BMC cardiovascular disorders*. 2012;12(1):1-6.
45. Bekler A, Barutçu A, Tenekecioglu E, Altun B, Gazi E, Temiz A, et al. The relationship between fragmented QRS complexes and SYNTAX and Gensini scores in patients with acute coronary syndrome. *Kardiologia Polska (Polish Heart Journal)*. 2015;73(4):246-54.
46. Stavileci B, Cimci M, Ikitimur B, Barman HA, Ozcan S, Ataoglu E, et al. Significance and usefulness of narrow fragmented QRS complex on 12-lead electrocardiogram in acute ST-segment elevation myocardial infarction for prediction of early mortality and morbidity. *Annals of Noninvasive Electrocardiology*. 2014;19(4):338-44.
47. Yanai H, Yoshida H. Secondary dyslipidemia: its treatments and association with atherosclerosis. *Global Health & Medicine*. 2021; 3(1):15-23
48. Caliskan B, Korkmaz A, Erdem F. Contribution of fragmented QRS on myocardial perfusion imaging in the assessment of functionally significant coronary artery stenoses. *European Review for Medical and Pharmacological Sciences*. 2016; 20: 1575-1581.
49. Bryer E, Stein E, Goldberg S. Multivessel Coronary Artery Disease: The Limitations of a "One-Size-Fits-All" Approach. *Mayo Clin Proc Inn Qual Out*. 2020;4(6):638-641
50. Kennedy HL, Goldberg RJ, Skizlo M. Acute Myocardial Infarction Revisited : A Community-Wide Study. *Clin. Cardiol*. 2011;2,455- 460.
51. Tanriverdi Z, Dursun H, Simsek MA, Unal B, Kozan O, Kaya D. The predictive value of fragmented QRS and QRS distortion for high-risk patients with STEMI and for the reperfusion success. *Annals of Noninvasive Electrocardiology*. 2015;20(6):578-85.

52. Ari H, Çetinkaya S, Ari S, Koca V, Bozat T. The prognostic significance of a fragmented QRS complex after primary percutaneous coronary intervention. Heart and vessels. 2012;27(1):20-8.
53. Michael MA, El Masry H, Khan BR, Das MK. Electrocardiographic signs of remote myocardial infarction. Progress in cardiovascular diseases. 2007;50(3):198-208.
54. Das MK, Saha C, El Masry H, Peng J, Dandamudi G, Mahenthiran J, et al. Fragmented QRS on a 12-lead ECG: a predictor of mortality and cardiac events in patients with coronary artery disease. Heart rhythm. 2007;4(11):1385-92.
55. Sorajja P, Gersh BJ, Cox DA, McLaughlin MG, Zimetbaum P, Costantini C, et al. Impact of multivessel disease on reperfusion success and clinical outcomes in patients undergoing primary percutaneous coronary intervention for acute myocardial infarction. European heart journal. 2007;28(14):1709- 16.
56. Lee JH, Park HS, Ryu HM, Lee H, Bae MH, Lee JH, et al. Impact of multivessel coronary disease with chronic total occlusion on one-year mortality in patients with acute myocardial infarction. Korean circulation journal. 2012;42(2):95-9.
57. Das MK, Zipes DP. Fragmented QRS: a predictor of mortality and sudden cardiac death. Heart rhythm. 2009;6(3):S8-S14.
58. Yesin M, Çağdaş M, Kalçık M, Rencüzoğulları I, Karabağ Y, Gürsoy M, et al. The relationship between fragmented QRS complexes and syntax II scores in patients with ST-segment elevation myocardial infarction. Elsevier Inc. 2018; 825–829.
59. Fedulaev Y , Makarova I, Pinchuk T. Fragmented QRS as a marker of coronary atherosclerosis severity evaluated by the Gensini score. Elsevier Masson SAS. 2021; 143–147.
60. Erdoğan T, Kocaman S, Çetin M, Çanga A, Durakoğlugil M, Çiçek Y, et al. Relationship of fragmented QRS complexes with inadequate coronary collaterals in patients with chronic total occlusion. J Cardiovasc Med. 2012;13(8):499-504
61. Eyuboglu M, Ekinci M, Karakoyun S, Kucuk U, Senarslan O, Akdeniz B. Fragmented QRS for Risk Stratification in Patients Undergoing First Diagnostic Coronary Angiography. Arq Bras Cardiol. 2016; 107(4):299-304.